

8/8/16

Reasoning & Aptitude

Gate

$$4Q \times 2\text{marks} = 8m$$

$$1Q \times 1m = 1m$$

E.S.E.

R/A 20 marks

✓ (2010-2013) → 9 marks

2014 (M.E.) → 12 marks

2015] → 10 marks
2016]

→ (20-25) %
P.S.U. (PSC + state engg. services)

C.S.A.T. → paper II

✓ 100 balls → 99 balls (10gms) each.
↳ 1 ball (9gms) faulty.

What is minimum no. of weighings required on a beam balance so as to find the faulty ball?

locate the faulty ball
our objective

Sol

B.B → 3^n

1 → 3^1 → 1

4 → 3^2 → 2

10 → 3^3 → 3

28 → 3^4 → 4


82 → 3^5 → 5 ✓ Ans.

✓ 100 Balls → 99 balls (10gms) each
 ↳ 1 ball (9gms) faulty

Minm. no. of weightage req. on a spring Balance.

Sol

always to ensure
 an answer keeping
 in mind the
worst case.



previous
 qn. (B.B.)

Spring Balance 2^n ✓

answer from this ball	Balls	→
1	2	2^1 → 1
3	4	2^2 → 2
5	8	2^3 → 3
9	16	→ 4
17	32	→ 5
33	64	→ 6
65	128	→ 7

✓ 10 Blue / 12 Grey
 ↳ Dark Room

Min (pair) → 3.

Min (Blue pair) → 14.

✓ Digital Balance → spring Balance.

CHAPTER 1

Number System

① Factors :- factors are the set of no.'s which will divide a given no. completely.

$$\begin{array}{r|l} 2 & 72 \\ \hline 2 & 36 \\ \hline 2 & 18 \\ \hline 3 & 9 \\ \hline 3 & 3 \\ \hline \end{array}$$

Factors \longleftrightarrow Divisors.

→ examiner denotation.

$$1. 72 = 2^3 \times 3^2 = \underline{4 \times 3} = 12 \text{ factors}$$

1, 2, 3, 4, 6, 8, 9, 12, 18, 24, 36, 72

$$\bullet 120 = 2^3 \times 3^1 \times 5^1 = \underline{4 \times 2 \times 2} = 16 \text{ factors}$$

1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120

$$\begin{array}{l} 2^0 \rightarrow 3^0 (1) \\ \rightarrow 3^1 (3) \\ \rightarrow 3^2 (9) \end{array}$$


$$\begin{array}{l} 2^1 \rightarrow 3^0 (2) \\ \rightarrow 3^1 (6) \\ \rightarrow 3^2 (18) \end{array}$$

$$\begin{array}{l} 2^2 \rightarrow 3^0 (4) \\ \rightarrow 3^1 (12) \\ \rightarrow 3^2 (36) \end{array}$$

$$\begin{array}{l} 2^3 \rightarrow 3^0 (8) \\ \rightarrow 3^1 (24) \\ \rightarrow 3^2 (72) \end{array}$$

Note :- $N = a^p \times b^q \times c^r$
Total factor = $(p+1)(q+1)(r+1)$

where a, b, c are distinct prime no.'s and p, q and r are natural no.'s.



2	10800
2	5400
2	2700
3	900
3	300
2	100
2	50
5	25
5	5

$(3) \times (5) \times (6) = 60$

$$\therefore N = 2^3 \times 3^2 \times 5^3$$

① Total factor (Tf) ($\sqrt{48}$)


② odd f ($\sqrt{12}$) $(3 \times 4) \checkmark$ ($2 \rightarrow \cancel{2}$)

③ even f $(48 - 12 = 36) \checkmark$

④ perfect square ($\sqrt{8}$) = $2 \times 2 \times 2 = 8$

⑤ perfect cubes ($\sqrt[3]{4}$) = $2 \times 1 \times 2 = 4$


Sol ① $4 \times 3 \times \cancel{4} = 48$



④ for perfect square, power should be multiply of $\underbrace{2}$ and $\textcircled{0}$.
even

$$2^4 \times 3^2$$
$$\cancel{2^2} \times 3^2$$

$$72 = 2^3 \times 3^2$$



$$2 \times 2 = 4$$

$$2^0 - 3^0 (1)$$
$$3^2 (9)$$

$$2^2 - 3^0 (4)$$
$$3^2 (36)$$

⑤ for no. to be perfect cube, power have to multiply of 3 and 0.

$$2^6 \times 3^3$$
$$\cancel{2^3} \times 3^3$$

Q How many factors of no. 72 are multiply of 6.

Sol

$$72 = 2^3 \times 3^2$$

$$\begin{array}{r|l} 6 & 72 \\ \hline & \end{array}$$

$$\begin{array}{l} (2 \times 3) (2^2 \times 3^1) \\ \swarrow \quad \searrow \\ 6 \quad \quad \quad 6 \\ \swarrow \quad \searrow \\ (1, 2, 3, 4, 6, 12) \end{array}$$

Ans

Q $120 = 2^3 \times 3^1 \times 5^1$

$$= 2^2 \times 3^1 (2^1 \times 5^1)$$

$2 \times 2 = 4$ Ans


$$= 12 (1, 2, 5, 10)$$

Q

$$\begin{array}{l} 30 \quad \quad \quad 18 \text{ } \checkmark \text{ } \underline{\underline{\text{Ans}}} \\ \uparrow \quad \quad \quad \swarrow \\ (2 \times 3 \times 5) \quad (3 \times 2 \times 3) \\ (2^2 \times 3^1 \times 5^2) \end{array}$$

* Prime factor :-

To
hell of
higher powers.



② Factorial :- is a product of 2 no.'s
 ↳ Multiplication of Natural NO. from 1 to N.

Q: $1! + 2! + 3! + 4! + 5! + 6! + 7! + \dots + 99!$ \Rightarrow unit digit

\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow
 1 2 6 24 0 0 0 0

Sol: $5! = 1 \times 2 \times 3 \times 4 \times 5 = 120$

$6! = 6 \times 5! = 720$

$7! = 7 \times 6 \times 5! = 5040$

first 4 no.'s

$$\begin{array}{r} 33 \\ +120 \\ +720 \\ \hline 0 \\ 0 \\ 0 \\ \hline 3 \end{array}$$

Ans $\rightarrow 3 \checkmark$

Note :- $5!$ onwards, every ! ends with atleast a single 0.

Q: $100!$ ends with how many 0?

Sol: $100! = 1 \times 2 \times 3 \times 4 \times 5 \times 6 \times \dots \times 99 \times 100$

$5 \rightarrow$ ~~on~~ $\frac{y}{e}$

$\frac{100}{5} = 20$ [5, 10, 15, 20, ..., 100] $\approx 5^1$

$\frac{20}{5} = 4$ [25, 50, 75, 100] $\approx 5^2$

24

Q $\frac{100}{3^n}$ (maximum power of 3 contained in $100!$)

Sol $100! = 1 \times 2 \times 3 \times 4 \times 5 \times 6 \times \dots \times 100$

$$\begin{aligned} \frac{100}{3} &= 33 \quad [3, 6, 9, 12, 15, \dots, 99] \approx 3^1 \\ &+ \\ \frac{33}{3} &= 11 \quad [9, 27, 36, \dots, 99] \approx 3^2 \\ &+ \\ \frac{11}{3} &= 3 \quad [27, 54, 81] \approx 3^3 \\ &+ \\ \frac{3}{3} &= 1 \quad [81] \approx 3^4 \\ &+ \\ &= 48 \quad \checkmark \end{aligned}$$

Q $\frac{100}{7^n}$

Sol $\frac{100}{7} = 14 \quad [7, 14, 21, \dots, 98] \approx 7^1$

$$\begin{aligned} \frac{14}{7} &= 2 \quad [49, 98] \approx 7^2 \\ &= 16 \quad \checkmark \end{aligned}$$

Q $\frac{100}{15^n}$

Sol $\frac{100}{15} = 6 \quad [15, 30, \dots, 90]$

$$100! = 1 \times 2 \times 3 \times 4 \times 5 \times \dots \times 99 \times 100$$

$$\frac{100!}{(3 \times 5)^n}$$

roti sabji

since limitation $\rightarrow 100! / 15^n \Rightarrow 24 \times 5$ times

$$100! = 3^48 \times 5^{24}$$

$$= (3 \times 5)^{24} \times (3)^{24}$$

$$= (15)^{24}$$

(15 → not prime no.
hence, बार बार अनेक बार)

Q A no. (of exact) has exactly 3 prime factors ($a^b \times b^c \times c^r$)
125 factors of the number are perfect squares.

Ans 27 factors of the number are — cube.

then overall Total factors of the no. are ?

Q Find the No. of trailing 0's.

(a) $1! \times 2^2 \times 3^3 \times \dots \times 100^{100}$

(b) $1! \times 2! \times 3! \times \dots \times 100!$

* BASE SYSTEM :-

$(25)_{10} = ((2^4 2^3 2^2 2^1 2^0)_{(2)})_{(2)}$

for Reaching to the Base (2).

2	25	Remainder
2	12	1 ↑
2	6	0
2	3	0
	1	1

$(16 * 8 + 0 + 0 + 1)$

$(25)_{10}$

MOHIT CHOUKSEY

$$\checkmark \begin{array}{r} \text{hr} \quad \text{Min.} \quad \text{Sec} \\ 3 \quad : \quad 24 \quad : \quad 36 \\ + 2 \quad : \quad 35 \quad : \quad 24 \\ \hline 6 \quad : \quad 0 \quad : \quad 0 \end{array} \quad b=60$$

if hr Min. Sec \rightarrow not given,
Then also Base 60 \checkmark .

\downarrow every time we are making Base 60 to zero.

$$\checkmark \begin{array}{r} \text{①} \quad \text{①} \\ \text{①} \quad 8 \quad 7 \quad 3 \\ + \quad 1 \quad 2 \quad 7 \\ \hline 10 \quad 0 \quad 0 \end{array} \quad \begin{array}{l} \text{Base} \\ \underline{10} \rightarrow \underline{0} \end{array}$$

$$\checkmark \begin{array}{r} \text{①} \quad \quad \quad \text{①} \\ \text{hr} \quad : \quad \text{Min.} \quad : \quad \text{Sec} \\ 3 \quad : \quad 24 \quad : \quad 36 \\ + 2 \quad : \quad 45 \quad : \quad 32 \\ \hline 6 \quad : \quad 10 \quad : \quad 8 \end{array} \quad b=60$$

$(60 \rightarrow 1 \rightarrow \text{①})$

$$\checkmark \begin{array}{r} 3 \quad : \quad 24 \quad : \quad 36 \\ 2 \quad : \quad 45 \quad : \quad 32 \\ \hline 6 \quad : \quad 10 \quad : \quad 8 \end{array}$$

$$36 + 32 = \overset{?}{\text{base}} + 8$$

$$\boxed{b=60} \checkmark$$

Q. (Gate)
2010
(2 marks)

$$\begin{array}{r}
 \textcircled{1} \quad 7+b = b+5 \rightarrow b=8 \\
 \textcircled{1} \\
 \begin{array}{r}
 1 \quad 3 \quad 7 \\
 2 \quad 7 \quad 6 \\
 \hline
 4 \quad 3 \quad 5 \\
 \hline
 \end{array}
 \end{array}$$

$b=8$

$$\begin{array}{r}
 \textcircled{1} \quad 1+2 = \\
 \textcircled{1} \\
 \begin{array}{r}
 7 \quad 3 \quad 1 \\
 + 6 \quad 7 \quad 2 \\
 \hline
 \textcircled{1} 6 \quad 2 \quad 3 \\
 \hline
 \end{array}
 \end{array}$$

$b=8$

$$7 + b = 8 + 5 \\
 \textcircled{13} \quad \downarrow \\
 \text{Base}$$

carry forward करके
when = or > base

Q. II.

$$\begin{array}{r}
 \begin{array}{r}
 7 \quad 6 \\
 3 \quad 2 \\
 1
 \end{array} \\
 \text{borrow} \\
 \begin{array}{r}
 (-) \quad 6 \quad 7 \quad 2 \\
 \hline
 0 \quad 3 \quad 7 \\
 \hline
 \end{array}
 \end{array}$$

$b=8$

(1-2) Not possible
hence Borrow करके

Q. III.

$$\begin{array}{r}
 \textcircled{1} \quad \textcircled{1} \quad \textcircled{1} \quad \textcircled{1} \\
 \begin{array}{r}
 4 \quad 4 \quad 4 \quad 2 \\
 + \quad 2 \quad 2 \quad 6 \\
 \hline
 1 \quad 0 \quad 0 \quad 1 \\
 \hline
 \end{array}
 \end{array}$$

alter

$$\begin{array}{r}
 \begin{array}{r}
 2 \quad 3 \quad 4 \quad 7 \\
 1 \quad 2 \quad 3 \quad 2
 \end{array} \\
 - \quad \begin{array}{r}
 1 \quad 6 \quad 5 \quad 6 \\
 \hline
 0 \quad 3 \quad 5 \quad 3 \\
 \hline
 \end{array}
 \end{array}$$

$b=7$

$$2 + 6 = b + 1 \\
 \textcircled{8} \quad \textcircled{b=7}$$

$2+b=8$ but $\frac{1}{7}$ more
 \downarrow
hence
7 is base

MOHIT CHOUKSEY

Q95
(Gate 2014)

$$(7 \ 5 \ 2 \ 6)_8 - (Y)_8 = (4364)_8$$

$$\begin{array}{r} (7 \ 5 \ 2 \ 6) \\ (-) 6 \\ \hline 4 \ 3 \ 6 \ 4 \end{array}_8$$

$$\begin{array}{r} \cancel{7} \ \cancel{5} \ 2 \ 6 \\ - \ 4 \ 3 \ 6 \ 4 \\ \hline 3 \ 1 \ 4 \ 2 \end{array}_8$$

$$6 + y_1 = 6 + 4$$

$$6 + y_2 = 8 + 4$$

$$6 + y_3 = 12$$

* $3 \longrightarrow 3, 6, 9, 12, \dots$
 $4 \longrightarrow 4, 8, 12, \dots$

$$k \times \text{LCM}(3 \times 4)$$

$$(12k)$$

* no. divisible by (2, 3, 5)

$$\hookrightarrow \text{LCM}(2, 3, 5) k$$

$$(30k)$$

MOHIT CHOUKSEY

118 Red light flashes \rightarrow $\left[\begin{array}{l} R(3 \text{ times} \rightarrow \text{every } 2 \text{ min}) \\ G(5 \text{ times} \rightarrow 3 \text{ min}) \end{array} \right] \rightarrow 120 \text{ sec.}$
 $\rightarrow 180 \text{ sec}$

$$\left(\begin{array}{l} R_1 \\ 40 \text{ sec} \end{array}, \begin{array}{l} G_1 \\ 36 \text{ sec} \end{array} \right)$$

$$\left(\begin{array}{l} R_1 \\ 40 \end{array}, \begin{array}{l} G_1 \\ 36 \end{array} \right) \text{ secs}$$

$$= 360 \text{ secs}$$

$$\approx 6 \text{ mins}$$

2	40, 36
2	20, 18
2	10, 9
5	5, 9
3	1, 9
	1, 3

$$\checkmark 1 \text{ hr} = \frac{60 \times 60}{360}$$

10 times

$$\text{LCM} \left(\frac{a}{b}, \frac{c}{d}, \frac{e}{f} \right) = \frac{\text{LCM}(a, c, e)}{\text{HCF}(b, d, f)}$$

Method alter

$$\checkmark \text{LCM} \left(\frac{2}{3}, \frac{3}{5} \right) \text{ min} = \left(\frac{6}{1} \right) \text{ mins}$$

$$\text{So within 1 hr} \rightarrow \frac{60 \text{ min}}{6} = 6 \text{ times}$$

\rightarrow if question says, they flash together at the beginning
 add '1' to the answer.



means $t = 0$
 $\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$

MOHIT CHOUKSEY

9/8/16

CYCLICITY

:- If a no. is ending in 2, its square have to end in 4, cube \rightarrow 8, quad \rightarrow 6.



Q. $(732)^{4(47)} \text{Re } 3 \dots \dots = (u =)$
 refer to cyclicity chart of 2

Ans $u = 8 \checkmark$

	unit place			
$4n+1 \rightarrow$	2	3	7	8
$4n+2 \rightarrow$	4	9	9	4
$4n+3 \rightarrow$	6	7	3	2
$4n \rightarrow$	8	1	1	6


4 \swarrow odd
 6 \swarrow even

9 \swarrow odd
 1 \swarrow even

[0, 1, 5, 6]

Q. $(74)^{91(\text{odd})}$ unit place = \dots (4)
 $(74)^{92(\text{even})}$ " " " " (6)

Q. $(237)^{254} \times (738)^{227} \times (76)^{2401} \times (79)^{5407}$



Pg 90 Gate 2016

Q 178

$21^{870} \times 146^{127} \times 3^{124}$

~~$1 \times 6 \times 8 \times 1$~~

~~7 ans~~

$1 + 6 \times 1 = 7$ ans

MOHIT CHOUKSEY

* REMAINDERS

any no. can be written in the form

$N = \text{Remainder} \pmod{\text{Divisor}}$

$\checkmark 80 = 8m(9)$
 \downarrow
 mod

$\checkmark 26 = 5m(7)$

$$x = y \pmod{m}$$

$$x - y = 0 \pmod{m}$$

Take $80 \leftarrow 8m(9)$

$$72 = 0 \pmod{9}$$

80 chocolates \rightarrow 9 students

$$80 = (-1)m(9)$$

$$80 + 1 = 0 \pmod{9}$$

$$81 = 0 \pmod{9}$$

$$26 \leftarrow -2m(7)$$

Rule-1 \rightarrow $\boxed{+, -, \times}$

$$a = b \pmod{c}$$

$$d = e \pmod{c}$$

$$f = g \pmod{c}$$

$$a \times d \times f = b \times e \times g \pmod{c}$$

$$b \times e \times g < c$$

$$\begin{array}{l} \text{if } a \\ +d \\ -f \end{array} \quad \begin{array}{l} b \\ +e \\ -g \end{array}$$

$$b + e - g < c$$

Q. Eg:-

$$1421 \times 1423 \times 1425$$

$$\begin{array}{l} 5 \times (-5) \quad 12 \times (-3) = 75 \\ 5 \times \quad \quad 7 \times \quad \quad 9 \times 12 \end{array}$$

Recd. Ans

$$\begin{array}{r} 1421 \times 1423 \times 1425 \\ \hline = 315 \\ \hline 12 \\ \hline = 3 \\ \text{Rem} \end{array}$$

$$1421 = 5m(12)$$

$$1423 = (-5)m(12)$$

$$1425 = (-3)m(12)$$

$$\begin{aligned} 1421 \times 1423 \times 1425 &= (-5)(-3)m(12) \\ &= 15m(12) \\ &= 3m(12) \end{aligned}$$

Rule-2

$$a = b \pmod{c}$$

$$a^n = b^n \pmod{c}$$

$$b^n < c$$

Q. $2^{600} \div 15$

$$24 = 1 \pmod{15}$$

$$(24)^{150} = (1)^{150} \pmod{15}$$

$$2^{600} = 1 \pmod{15}$$

MOHIT CHOUKSEY

Pg 40
Q6 and Q7

Sol
$$\begin{array}{r} 6 \\ 49 \\ \times 7 \\ \hline 343 \end{array}$$

$784 \div 342$

$$\underbrace{7^3}_a = \underbrace{(1)}_b \pmod{\underbrace{(342)}_c}$$

$$(7^3)^{28} = (1)^{28} \pmod{(342)}$$

 ✓

Q7 $(15^{23} + 23^{23}) \div (19)$

$$\begin{array}{r} (15)^{23} = (-4)^{23} \pmod{19} \\ + (23)^{23} = (+4)^{23} \pmod{19} \\ \hline 15^{23} + 23^{23} = 0 \pmod{19} \end{array}$$

Q (a)
$$\frac{10^{10} + 10^{100} + 10^{1000} - 10^{10000}}{3}$$

Sol

$$\begin{aligned} (10)^{10} &= (1)^{10} \pmod{3} \\ + (10)^{100} &= (1)^{100} \pmod{3} \\ + (10)^{1000} &= (1)^{1000} \pmod{3} \\ + (10)^{10000} &= (1)^{10000} \pmod{3} \\ \hline &= 2 + 1 - 1 \\ &= 2 \checkmark \end{aligned}$$

(b) $5^{625} \div 7$

$$\begin{aligned} 5^5 &= 3 \pmod{7} \\ (5)^{125} &= (3)^{125} \pmod{7} \\ 3^{125} \div 7 & \\ (3^5)^{25} \div (5) \pmod{7} & \end{aligned}$$

$$\begin{array}{r} 25 \\ \times 5 \\ \hline 125 \\ \times 5 \\ \hline 625 \\ \times 5 \\ \hline 3125 \\ \hline 446 \\ 7 \overline{) 3125} \\ 28 \downarrow \\ \hline 32 \\ 28 \downarrow \\ \hline 45 \\ 42 \downarrow \\ \hline 3 \end{array}$$

$$\begin{array}{r} 9 \\ \times 3 \\ \hline 27 \\ \times 3 \\ \hline 81 \\ \times 3 \\ \hline 243 \\ 37 \\ 7 \overline{) 243} \\ -21 \\ \hline 33 \\ -28 \\ \hline 5 \end{array}$$

SIR $5^{625} \div 7 \Rightarrow 5^3 = 6 \pmod{7}$

also

$5^3 = (-1) \pmod{7}$

$(5^3)^{208} = (-1)^{208} \pmod{7}$

$$\begin{aligned} 5^{624} &= 1 \pmod{7} \\ \times (5)^1 &= (-2) \pmod{7} \text{ or } 5 \end{aligned}$$

$$\begin{array}{l} (-2) \pmod{7} \\ \hline 5 \pmod{7} \end{array}$$

$$\begin{array}{r} 5 \leq 7 \\ 0 \\ 7 \overline{) 5} \\ -0 \\ \hline 5 \end{array}$$

MOHIT CHOUKSEY

$$* (5)^{625} = (-2)^{625} \pmod{7}$$

\downarrow
 No need
 any no.

hence, Taken smaller power.

$$5^3 = (-2)^3 \pmod{7}$$

$$5^3 = -8 \pmod{7}$$

$$5^3 = -\frac{8}{\sqrt{8}} \pmod{7}$$

aqaya \rightarrow 8
 341 \rightarrow 8

$$\sqrt{(5^2)^3} = (4)^3 \pmod{7}$$

$$(5^6)^{104} = (1)^{104} \pmod{7}$$

$\frac{64}{7} =$
 $Re \rightarrow 1$

$$5^{624} = 1 \pmod{7}$$

MOHIT CHOUKSEY

Time and Work Calendar

36 365 d, 5 hrs, 48 mins, 11 secs - - - - -

$365d \approx 6 \text{ hrs}$

- ① Every multiple of 4 is a Leap year (4, 8, 12, 16 - - - - LY)
- ② Century year is Non Leap year (100, 200, 300, - - - - NLY)
- ③ Every 4th century year is LY (400, 800, 1200, - - - - LY)

ordinary year
 $1(O.Y.) = 365d = \frac{52 \times 7}{1} + 1 \text{ odd day}$

$1(O.Y.) = \left(\frac{365}{7}\right)d$ Remainder (Re) 1 odd day.

$1(L.Y.) = \left(\frac{366}{7}\right)d$ Re 2 odd day.

within 1st 100Y \longrightarrow $24(L.Y.) + 76(O.Y.)$
 $\times 2(Re) + \times 1(Re)$

$48 + 76 = \frac{124}{7} \text{ Re } 5 \text{ odd day}$

since never
 Re can't be
 more than divisor
 or no of odd days
 can't be more
 than 7
 hence \div by 7 to
 get
 (5)
 odd day.

4) $\times 100 Y \rightarrow 5$ odd days

$\times 200 Y \rightarrow 3$ odd days

$\times 300 Y \rightarrow 1$ —

$\checkmark 400 Y \rightarrow 6 \rightarrow \dots + 1 = \frac{7}{7} = 0 \text{ Re}$
 \downarrow
0 odd days

$\frac{00}{00} \times 4 \uparrow$
hence
extra
1 day

5) 1st odd day is MONday

(Gregorian calendar \rightarrow 01 01 AD)

Q 28th Aug 1994
1900 \rightarrow 1

?

6) (a) 400 \rightarrow 0

1600 \rightarrow 0

300 \rightarrow 1

1900 \rightarrow 1

Sol 0 - 1900 \rightarrow 1

1900 - 93 \rightarrow 4

94 \rightarrow 2

7

$Y = 93 = \begin{matrix} L.Y. & O.Y. \\ 23 & + & 70 \\ \times 2 & + & \times 1 \end{matrix}$

$46 + 70 = \frac{116}{7} \text{ Re } (4) \checkmark \text{ odd day}$

1994

J ~~31~~ \leftarrow do well
31 $\frac{7}{7}$ Re (3) \downarrow odd
F 29 1 \rightarrow 94 is not a leap year
F 0 M 3 A 2 My 3 J 2 3 Aug 0

Q. 9th Aug 2016

<G>(b) $\begin{matrix} (\because 400 \rightarrow 0) \\ 0 - 2000 \rightarrow 0 \end{matrix}$

Sol

2000 \rightarrow 0
 15 \rightarrow 4
 16 \rightarrow 5

 9

Y = L.Y. + O.Y.
 15 = 3 + 12

$\begin{matrix} \times 2 & \times 1 \\ \hline 6 + 12 = \frac{18}{7} \end{matrix}$ (Re 4)

2016
 Leap year

J F M A M J J A
~~3~~ ~~1~~ ~~3~~ 2 3 2 3 2

$\frac{9}{7}$ Re (2) \rightarrow Tuesday

$\frac{9}{7}$ (Re 2)

Q If 15th Aug 1947 was Friday, then 26th January 1950 was _____.

Sol 26th Jan. 1950

0 - 1900 \rightarrow 1

LY NLY
 49 \rightarrow 12 37
 $\times 2 \quad \times 1$

~~48 + 27 = 75~~
~~7~~

24 + 37 = 61

$\frac{61}{7} =$

$\begin{matrix} (12) & (48) \\ \hline 12 \\ -50 \\ \hline 4 \end{matrix}$
 $\begin{matrix} 7 \overline{) 75} \\ -7 \downarrow \\ \hline 65 \end{matrix}$

MOHIT CHOUKSEY

812

26th Jan 1950

Paper print → Sunday
1st odd day → Monday

1900 → 1	49	LY	0Y
49 → 5	12		37
50 → 5		x 2	x 1
		<hr/>	
		24 + 37	= 61
			7

(11/7) Re 4 (61/7) Re 5

1950

Thursday

26 Jan Re 5
7


Alternate Method

Let 15th August 1947 = Fri = 0th odd day

	Aug (16/7)	Sep	Oct	Nov.	Dec		
1948	2	2	3	2	3		
1948	2	← 366 (L.Y.)				2	2
1949	1	← 365 (O.Y.)				+2	3
1950	5	← 26th Jan.				+3	2
		7				<hr/>	<hr/>
						7	7

6 → Thursday

MOHIT CHOUKSEY



$$n(A \cup B) = + [n(A) + n(B)] - [n(A \cap B)]$$

$$n(A \cup B \cup C) = + [n(A) + n(B) + n(C)] - [n(A \cap B) + n(B \cap C) + n(A \cap C)] + [n(A \cap B \cap C)]$$


$n(A \cup B \cup C \cup D)$
 $\wedge \vee D$

$+ \sum n(A)$	$\rightarrow 4 \text{ values}$	$4C_2$	AB		ABC
$- \sum n(A \cap B)$	$\rightarrow 6 \text{ values}$	$4C_2$	AC		ACD
$+ \sum n(A \cap B \cap C)$	$\rightarrow 4 \text{ values}$	$4C_3$	AD		ABD
$- n(A \cap B \cap C \cap D)$	$\rightarrow 1 \text{ value}$	$4C_4$	BC		BCD
			BD		
			CD		

MOHIT CHOUKSEY

C → Cricket
 B → Basketball
 H → Hockey

C & H → outside → 30
 C & H only → inside → 20
 C & H but not B → C & H only
 ↓
 20



Q How many students are playing any of these 3 games
 (or) atleast one of 3 games

$$n(A \cup B \cup C) = [100 + (x + y + z)]$$


let → x, y, z ← naming

Q none of these 3 games

$$\leftarrow T - n(A \cup B \cup C)$$

 Total

MOHIT CHOUKSEY



(a) atleast 2 of games = (B) + (B)

(a) atleast 1 of games = (R) + (B) + (B)

= sum of all the values.

Cricket only → (inside)


Cricket → (बाहर)

1,2,3 Pg 48

a (20%) ✓


b (10%) ✓

c $20 + 0 + 10 = 30\%$ ✓



Pg 48 8 to 11



- 80 ✓
- 170 ✓
- 150 ✓
- 30 ✓




$$\begin{array}{r} 220 \\ + 50 \\ + 80 \\ \hline 350 \end{array}$$

$$\begin{array}{r} 500 \\ - 350 \\ \hline 150 \end{array}$$

$$\begin{array}{r} 200 \\ - 80 \\ \hline 120 \\ + 30 \\ \hline 150 \\ + 40 \\ \hline 190 \\ + 50 \\ \hline 240 \end{array}$$





Sol



MOHIT CHOUKSEY

Q 166



$$T - 30 = n(A \cup B) = 120$$

$$n(A) + n(B) - n(A \cap B) = 120$$


$$55 + 85 - \text{---} = 120$$

$$n(A \cap B) = 20$$

MOHIT CHOUKSEY

Pg 68

Q1



Pg 79
Q 13 to 16

→ logical Venn Diagram → eyesight Test.

Prepp
Your Personal Exam Guide

MOHIT CHOUKSEY

10/8/16

Q9 $S_2 = 1 \times 2 + 2 \times 3$

$S_2 = 8$

put $n = 2$ ops

$S_2 = 8 (c)$

$\sum T_n = \sum n(n+1)$

$S_n = (\sum n^2 + \sum n)$

Q T1 $\Rightarrow A = 2^{172} - 2^{171}$

$A = 2^{171} (2 - 1)$

$A = 2^{171} =$

Q Ans. N lies b/w $9 < N < 1000$

$S_N + P_N = N$

Q $(\underbrace{1\ 2\ 1\ 2\ 12\ 12\ 12\ \dots\ 12}_{99}) \left(\begin{matrix} (12 \text{ written}) \\ (150 \text{ times}) \end{matrix} \right)$ calc. Remainder
(300 digit no.)

Q n is a 3^{digit} natural no. on the base of 10 and converted into base of 7 and base 9, how many such no's are there.

$(a\ b\ c)_7$ $(c\ b\ a)_9$

digits reverse ho jati hai

TIME & WORK

A \rightarrow 16 d

1 day work of A $\rightarrow \frac{1}{16}$

13 day work of A $\rightarrow \frac{13}{16}$

$$\begin{aligned} \text{Left over work} &= 1 - \frac{13}{16} \\ &= \frac{3}{16} \end{aligned}$$

$$\frac{1}{2} \times 10 \xrightarrow{\text{A}} = 2B \xrightarrow{\text{10 days}}$$

\downarrow
5 days

$$A = \frac{1}{2} B \xrightarrow{\text{10 days}}$$

\downarrow
 $2 \times 10 = 20 \text{ days}$

Q A is 4 times as eff. as B and takes 15 days less than B to finish a work. In how many days will the work get finished/done if A and B are working together

Sol

$$\frac{1}{4}(4x) \quad \text{A} = 4B \xrightarrow{\text{4x days}}$$

x days

$$3x = 15$$

$$\boxed{x = 5}$$

A \rightarrow 5 days \rightarrow one day work $\rightarrow \frac{1}{5}$
 B \rightarrow 20 days \rightarrow " " " " $\rightarrow \frac{1}{20}$

$$\boxed{\left[\frac{1}{5} + \frac{1}{20} \right] = \left[\frac{1}{4} \right]}$$

in one day \leftarrow , $\frac{1}{4}$ th of work is completed

so 4 days

Ans

MOHIT CHOUKSEY

Alternate work concept

alone \rightarrow A = 12 days
 \rightarrow B = 16 days

Q In how many days will the work be done if A and B are working alternatively, beginning with A.

Sol

$$\begin{array}{l} \text{1st day of A} \\ \text{2nd day of B} \end{array} = \left[\left(\frac{1}{12} + \frac{1}{16} \right) \right] = \frac{7}{48} \times 6 = \frac{42}{48} = \frac{7}{8}$$

2 day work \times 6 cycles

$$\begin{array}{l} \text{12 days work} \\ \text{work} \end{array} = \frac{7}{8} \quad \parallel \quad (\text{LOW}) = \frac{1}{8}$$

on 13th day (A)

$$\frac{1}{8} - \frac{1}{12} = \frac{1}{24} \quad (\text{LOW})$$

on 14th day (B)

$$\frac{\frac{1}{24}}{\frac{1}{16}} = \frac{2}{3}$$

$13\frac{2}{3}$
A starts

Q if B starts the work.

Sol 12 days work = $\frac{7}{8}$, LOW = $\frac{1}{8}$

on 13th day (B)

MOHIT CHOUKSEY

Siv

1st day of B 2nd day of A

$$\frac{2 \text{ dw}}{\times 6} = \left[\frac{1}{16} + \frac{1}{12} \right] = \frac{7}{48} \times 6 = \frac{42}{48} = \frac{7}{8}$$

$$\frac{12 \text{ dw}}{=} = \frac{7}{8} \quad | \quad \text{LOW} = \frac{1}{8}$$

on 13th day
(B)

$$\frac{1}{8} - \frac{1}{16} = \frac{1}{16} \text{ [LOW]}$$


on 14th day
(A)

$$\frac{\frac{1}{16}}{\frac{1}{12}} = \frac{3}{4}$$

→ 13³/₄ days Ans
if B starts

Pg 47
Q 6

- A → 20
- B → 15
- C → 12



SIR

1st day work of (A & B)

$$\left[\frac{1}{20} + \frac{1}{15} \right] = \frac{7}{60}$$

(3/60) (4/60)

2dw

2nd day work of (A & C)

$$\left[\frac{1}{20} + \frac{1}{12} \right] = \frac{8}{60}$$

(3/60) (5/60)

7/60 + 8/60 = 15/60 = 1/4

2dw = 1/4

2dw × 4 = 1

8 dw = 1

2 days → 1/4th work

Q A = 10 days
 alone → B = 12 days
 ↘ C = 15 days

maxm. 2 people are allowed to work in any single day
 with no two consecutive day having same pair of people repeating

Then the minimum no. of days in which work can be done?

Sol

1st dw of (A and B) $\left(\frac{1}{10} + \frac{1}{12} \right) = \frac{11}{60}$ (most efficient + lesser efficient)

2nd dw of (A & C) $\left(\frac{1}{10} + \frac{1}{15} \right) = \frac{10}{60}$

2nd dw = $\frac{21}{60}$
 X 2 = $\frac{42}{60}$

4th dw = $\frac{42}{60}$ | low = $\frac{18}{60}$

on 5th day
 A & B $\frac{18}{60} - \frac{11}{60} = \frac{7}{60}$ (Low)

on 6th day
 A & C $\frac{7/60}{1/60} = 7/10$

5 $\frac{7}{10}$ days
 ✓ Ans

* Men-days Concept

Inversely proportional

$\uparrow a \propto \frac{1}{b} \downarrow$

$a = \frac{k}{b}$

$a \times b = k$

$a_1 \times b_1 = a_2 \times b_2$

$\uparrow m \propto \frac{1}{d} \downarrow$

$m \times d = k$

$m_1 \times d_1 = m_2 \times d_2$

if $(200 m \times 10 \text{ days}) = \underline{2000 m d}$

$$\begin{array}{ccc} 90m & \longrightarrow & 270d \\ 30m & \longrightarrow & x \end{array} \quad \begin{array}{l} \downarrow \\ 3 \text{ times} \\ \downarrow \end{array}$$

$\frac{1}{3} \text{rd}$

$$90 \times 270 = 30 \times x$$

$$\underline{x = 810}$$

Q5

$$(4m + 3w)^2 = (6m + 9w) \times 4$$

$$(8m + 6w) = (6m + 9w)$$

$$\boxed{2m = 3w}$$

$$\boxed{1m = 1.5w}$$

$$(20m + 6w)x = (6m + 9w) \times 4$$

$$(30w + 6w)x = (9w + 9w) \times 4$$

$$(36w)x = (18w \times 4)$$

$$\boxed{x = 2 \text{ days}}$$

Q11

$$\cancel{(5M + 7B) \times 24} =$$

$$(9m + 18b) \times 15 \times 8 = \sqrt{(3m + 6b)} \times x \times 8$$

$$3(3m + 6b) \times 15 = (3m + 6b) x$$

$$\boxed{x = 45 \text{ days}}$$

MOHIT CHOUKSEY

Q8

$$A = 15 \times 8 = 120 \text{ hrs}$$

$$B = 6\frac{2}{3} \times 9 = 60 \text{ hrs}$$

$$10 \left[\frac{1}{120} + \frac{1}{60} \right] x = 1$$

$$10 \left[\frac{1}{24} \right] x = 1$$

$$x = 4$$

Q9

$$A = 24 \text{ days}$$

$$B = 36 \text{ days}$$

~~$$\left(\frac{1}{24}\right)x + \left(\frac{1}{36}\right)x = 6$$~~

$$\left(\frac{1}{24}\right)x + 6 \left[\frac{1}{24} + \frac{1}{36} \right] = 1$$


$$\frac{x}{24} = \frac{7}{12}$$

$$x = 14 \text{ days}$$

Q

$$\left[\frac{1}{10} + \frac{1}{15} \right] = \frac{1}{6}$$


6 hrs



$$\left[\frac{1}{10} - \frac{1}{15} \right] = \frac{1}{30}$$

drainage pipe

30 hrs




MOHIT CHOUKSEY

Q

~~$$\left[\frac{1}{15} - \frac{1}{60} \right] = \frac{3}{60}$$

$$= \frac{1}{20}$$~~



SIR

$x \rightarrow$ mins

$$15 \left[\frac{1}{x} - \frac{1}{60} \right] = 1$$

↓
1 min work


Pq77
Q71

~~$\frac{1}{30}$~~

~~$$10 \left[\frac{1}{x} - \frac{1}{30} \right] = 1$$~~

~~$$\frac{1}{x} = \frac{1}{10} + \frac{1}{30}$$~~

~~$$\frac{1}{x} = \frac{3+1}{300}$$~~




SIR

$x \rightarrow$ mins

$$10 \left[\frac{1}{x} - \frac{1}{60} \right] = \frac{1}{2}$$

$$\frac{1}{x} - \frac{1}{60} = \frac{1}{20}$$

$$\frac{1}{x} = \frac{1}{20} + \frac{1}{60} =$$



MOHIT CHOUKSEY

Q144

	P	Q	R	S
Q	→	25	x	12
R	→	50	x	12
Q	→	(x-2)	x	12
R	→		x	18

Previous Qn
 $y \rightarrow \text{min}$
 $10 \left[\frac{1}{y} - \frac{1}{30} \right] = 1$
 half the tank
 $y = 7.5 \text{ min}$

$$\begin{array}{r} 25 \\ \underline{12} \\ 50 \\ \underline{25x} \\ 300 \end{array}$$

SIR Q = 25 x 12 = 300 hrs.

R 50 x 12 = 600 hrs

$\frac{1}{300} \rightarrow$ 1hr work of Q

$\frac{1}{600} \leftarrow$ 1hr work of R

$\left(\frac{5 \times 12}{300} \right) \rightarrow$ 1/5th of work

$$\frac{60}{300}$$

$\left(\frac{18 \times 7}{600} \right) \rightarrow$ own fraction of work

$$\frac{126}{600}$$

$\frac{60}{300} : \frac{63}{300}$

20 : 21 ✓

Q173
G2016

A → 6 hr A 1/6
 B → 4 hr B 1/4

$\frac{1}{6}$

SIR

A 3 $\left(\frac{1}{6} \right)$ = $\frac{1}{2}$ Re A = $\frac{1}{2}$
 B 3 $\left(\frac{1}{4} \right)$ = $\frac{3}{4}$ Re B = $\frac{1}{4}$

Pg 24

A = 3 x 4 = 12 Re A = 12
 B = 3 x 6 = 18 Re B = 6

$\left[1 - x \left(\frac{1}{6} \right) \right] = 2 \left[1 - x \left(\frac{1}{4} \right) \right]$ $x = 3$

**MOHIT
CHOUKSEY**

Pg 47
Q12

~~m~~

$$m + B = 160 \text{ Rs}$$

$$m = 3B$$

$$3B + B = 160 \text{ Rs}$$

$$B = 40 \text{ Rs}$$


$$m = 3 \times 40 = 120 \text{ Rs}$$

Q2

$$4m \times 40 = 7W \times 40$$


$$4m = 7W$$

Mixture



MOHIT CHOUKSEY

Q




①
$$\frac{Q_c}{Q_D} = \frac{P_D - MP}{MP - P_c}$$

Labels: Q_c (quantity cheaper), Q_D (quantity dearer), P_D (dearer price), MP (mean price), P_c (cheaper price).
 Note: "awara formula" (अवरा सूत्र)

$$\frac{Q_{20}}{Q_{30}} = \frac{30 - 24}{24 - 20} = \frac{6}{4} = \frac{3}{2}$$

$$\frac{15}{10} = \frac{15}{1} = \frac{12}{8} = \dots$$


Q



$$\frac{Q_I}{Q_{II}} = \frac{84 - 72}{72 - 62} = \frac{12}{10} = \frac{6}{5}$$

Labels: 6L ✓, 5L ✓

Q




$$10,000 \rightarrow 750$$

$$100 \rightarrow 7.5\%$$

MOHIT CHOUKSEY

③ Replacements formulae / Replacements.

Volm = k
conserve



Quantity of milk left after n^{th} operation

Initial ~~limited~~ quantity of milk [I.Q.] = $\left[\frac{a-b}{a}\right]^n = \left[1 - \frac{b}{a}\right]^n$

Qu. of Milk left after n^{th} opr = $IQ \times \left[1 - \frac{b}{a}\right]^n$

where a is initial quantity, b is quantity taken out everytime & replaced by water, n = no. of replacements/operations.

$$\left[x - \frac{10}{100} x \right] \quad x [1 - 0.1]$$


↑	↓
1.10x	0.90x
1.20x	0.80x
1.30x	0.77x
1.23x	

Quantity ----- of milk after 1st opr = $100 \left[1 - \frac{10}{100} \right]^1 = 100 \times 0.9 = 90 \checkmark$

----- 2nd " = ----- $= 100 \times 0.9 \times 0.9 = 81 \checkmark$

----- 3rd ----- $= 72.9$ and so on

like



Now

$$\text{-----} \text{-----} \text{-----} \text{ 2nd opr} = 100 \left[1 - \frac{10}{100} \right] \left[1 - \frac{20}{100} \right]$$

$$= 100 \times 0.9 \times 0.8$$

$$= 72\%$$

$$\text{Q Milk left} = 40 \left[1 - \frac{4}{40} \right] \left[1 - \frac{5}{40} \right] \left[1 - \frac{6}{40} \right]$$

$$= \textcircled{A}$$

$$\text{water left} = 40 - \textcircled{A}$$

Pg 69
Q7

$$10 \left[1 - \frac{1}{10} \right]^3 = 7.29$$

1 is 10% of 10.

Prepp
Your Personal Exam Guide

MOHIT CHOUKSEY

11/8/16
T4
Pg 48

$$\left(\frac{1}{A} + \frac{1}{B} = \frac{1}{12}\right) \left(\frac{1}{B} + \frac{1}{C} = \frac{1}{16}\right) \rightarrow B = 48$$

$$\frac{5}{A} + \frac{7}{B} + \frac{13}{C} = 1$$

$$5\left[\frac{1}{A} + \frac{1}{B}\right] + 2\left[\frac{1}{B} + \frac{1}{C}\right] + \frac{11}{C} = 1$$

$$5\left(\frac{1}{12}\right) + 2\left(\frac{1}{16}\right) + \frac{11}{C} = 1$$

$$C = 24$$

T5 → 48

PERCENTAGE

Q → A's salary is 20% more than that of B. By how much % is B's salary less than that of A.

Sol

$$B = 100, \quad A = 120 \quad \begin{array}{l} \text{20\% } \uparrow \\ \frac{-20}{120} = -\frac{1}{6} \approx \underline{16.6\%} \downarrow \end{array}$$

$$\begin{array}{l} \text{let } 100 \xrightarrow{10\% \uparrow} 110 \\ \frac{-10}{110} = -\frac{1}{11} \approx 9.09 \downarrow \end{array}$$

Q → A's salary is 20% less than that of B. By how much % is B's salary more than that of A.

$$B = 100, \quad A = 80 \quad \begin{array}{l} \text{20\% } \downarrow \end{array}$$


$$\frac{+20}{80} = \frac{1}{4} \approx \underline{25\%} \uparrow$$

$$\text{let } B = 100 \xrightarrow{\text{25\% } \downarrow} 75$$

$$\frac{+25}{75} = \frac{1}{3} \approx \underline{33.\bar{3}} \uparrow$$

$$\frac{20}{80} = \frac{1}{4}$$

$$25\%$$



* $R = a \times b$

a changes by $x\%$

b changes by $y\%$

$$\Delta R = x + y + \frac{xy}{100}$$

MOHIT CHOUKSEY

Ex:- $A = l \times b$

$D = a \times t$

Revenue (R) = Price of car (P) \times (N) No. of car

* $l = 20 \uparrow$ $b = 10 \uparrow$ 32%

$A = 1l \times 1b$

$A' = 1.2l \times 1.1b$

$A' = 1.32 lb$

or

$$20 + 10 + \frac{20 \times 10}{100} = 32\%$$

$l = 20 \uparrow$ $b = 10 \downarrow$

$A = l \times b$

$A' = 1.2l \times 0.9b$ 8%

$A' = 1.08 lb$

or

for every \uparrow \downarrow

$$20 - 10 + \frac{20(-10)}{100} = 8\%$$

every for decrease

*** ** PROFIT (%) LOSS

$$P = (SP - CP)$$

✓ selling price (SP)

✓ cost price (CP)

$$P\% = \left[\frac{(SP - CP)}{CP} \right] \times 100$$
$$L\% = \left[\frac{(CP - SP)}{CP} \right] \times 100$$

20% Profit \rightarrow $SP = CP \times 1.2$

\hookrightarrow SP is 20% above the cost price.

20% Loss \rightarrow $SP = CP \times 0.8$

\hookrightarrow SP is 20% below the cost price.

Q eggs are bought at the rate of 7 eggs for Rs. 1. If the shopkeeper wants to make a profit of 40%, how many eggs should he sell for 1 Rs.

Sol

$$CP \text{ (1 egg)} = \left(\frac{1}{7} \right)$$
$$SP \text{ of 1 egg} = \left(\frac{1}{7} \right) \times 1.4$$
$$=$$


MOHIT CHOUKSEY

Q ① A dishonest shopkeeper uses a false weight of 800 gm instead of 1 kg weight. If he promises to sell the goods at the cost price, then his profit %.

Q ② On selling 36 mangoes, a shopkeeper recovers a CP of 33 mangoes only. Find loss %

Sol 1

$$P = \frac{SP - CP}{CP}$$



SIR

$$\frac{\text{Profit}}{CP} = \frac{(\text{CP of } 200 \text{ gms})}{(\text{CP of } 800 \text{ gms})} = \frac{1}{4} \approx 25\%$$

Sol 2

$$\frac{L}{CP} = \frac{(\text{CP of } 3 \text{ Mangoes})}{(\text{CP of } 36 \text{ mangoes})} = \frac{1}{12} \approx 8.\bar{3}\%$$

Q 1) A dishonest milkman uses a false measuring vessel of 800 ml instead of 1 L and further adulterates milk with 20% water (free of cost). If he promises to sell the milk at the CP then his Profit %.

Sol

~~$$P = \frac{SP - CP}{CP}$$

$$P\% = \frac{CP \text{ of } 2}{CP \text{ of } 2}$$~~

1st cheating $\rightarrow \frac{CP \text{ of } 200 \text{ ml}}{CP \text{ of } 800 \text{ ml}} = \frac{1}{4} \approx 25\%$

1st cheating (without mixing water) \rightarrow 25%

$$(1.25) \times (1.2) = 1.50$$

or

$$25 \times 20 + \frac{25 \times 20}{100} = 50\%$$

ex

1 ml = 1 Rs

1000 ml ^m = 1000 Rs

(+) 200 ml _w = 0 Rs

1200 ml = 1000 Rs = Total cost Price (TCP)

$\frac{1}{6} = 16.6\% \text{ w}$

800 ml \rightarrow Rs. 1000

400 ml \rightarrow Rs. 500

1200 ml \rightarrow TCP = Rs 1500

$$\frac{P}{CP} = \frac{500}{1000} \times 100 = 50\%$$

other ex

1 ml = $\frac{1}{6}$ 1 Rs

800 ml ^m \rightarrow Rs 800

160 ml _w \rightarrow Rs 0

960 ml \rightarrow Rs. 800 = TCP

adulterate \rightarrow add.
puts \rightarrow
mixture does not
contain 20%
water

800 ml \rightarrow Rs 1000

160 ml \rightarrow Rs. 200

960 ml \rightarrow 1200 Rs

$$\frac{P}{CP} = \frac{400}{800} \times 100 = 50\%$$

beaker
2/21 2/21

beaker
1/5 m

MOHIT CHOUKSEY

Q 29 (Pg 72)

SP \rightarrow Rs 60/unit

TC_{Pris} = Rs 100

TP_{%fit} = Rs 500

SP = 600

\uparrow 30%

130
TC' = 130

₹ 60 ——— unit

100 ₹
Six

SP/unit = 60 Rs

Total CP = 100 Rs

Profit = 500 Rs

Profit % = $\frac{P}{CP} \times 100 = \frac{500}{100} \times 100 = 500\%$

~~SP = CP + Profit~~

~~SP = 600~~ CP' = 130 Rs

P.% = 500% same

Profit' = 650 Rs (+130)

SP' = 780

SP'/unit = $\frac{780}{10} = 78$


60×1.3 30%

lay men
dont
follow
this

Now
SIR

$$\begin{aligned}
 SP &= CP \times 1.0P \\
 SP \times 1.3 &= \underbrace{CP \times 1.3}_{CP'} \times 1.0P \\
 SP' &= CP'
 \end{aligned}$$

Profit doesnt depends
on no. of
units.



milkman

$$1.25 \times 1.20$$

$$25 + 20 \times \frac{25 \times 20}{100}$$

$$= 50\%$$

successive profit condⁿ → dhamaake pay dhamaaka.

$$\frac{-80 - 40 + (-80)(-40)}{100} = -88 \text{ off}$$

$$-120 + 32$$

$$100$$

$$\downarrow \times 0.9$$

$$90$$

$$\downarrow \times 0.9$$

$$81$$

$$\downarrow \times 0.9$$

$$72.9$$

Q

$(70\% + 30\%) \text{ off}$


$100 \times 0.3 \times 0.7 = 21$

$\text{if } \neq 10\%$

$\times 0.9 =$

-79%


MARKED PRICE → list price, labelled price, print price, MRP



[N.P]

MOHIT CHOUKSEY



marked price \rightarrow selling price
 and CP \rightarrow Profit.



Q. \rightarrow After offering a discount of 20%, a shopkeeper still manages to make a profit of 25%. By how much % is the mark price above the cost price.

Sol

$$MP \times 0.8 = SP = CP \times 1.25$$



$$MP = CP \times \frac{1.25}{.8}$$

$$MP = CP \times 1.5625$$

56.25%


✓

$$MP \times 0.8 = SP$$

$$MP \times \frac{4}{5} = SP$$

$$MP = SP \times \frac{5}{4}$$

$$MP = \frac{SP \times 1.25}{1}$$



$$25 + 25 + \frac{25 \times 25}{100}$$

56.25%

MOHIT CHOUKSEY

Two Rules

↳ Rule (1) Two articles are sold at a common SP (selling price) of Rs S each. one is sold at a profit of P% and another at a loss of P%, then effectively there is always a loss during the entire transaction

$$\text{(value)} \quad \boxed{\text{Loss} = \frac{2P^2 S}{(100^2 - P^2)}} \quad (\text{Rs})$$

$$\boxed{\text{Loss}\% = \frac{P^2}{100} \%}$$

↳ Rule (2) Two articles are bought at a common CP, one is sold at a profit of P% and another at a loss of P%, then effectively there is no profit no loss.

Q Two shirts are sold at a common SP of Rs 480 each, 1 is sold at a profit of 20% and ~~20~~ another at a loss of 20%. then find loss and loss %.

Sol

$$\begin{array}{l} SP_1 = SP_2 = ₹ (480) \text{ each} \\ SP_1 = CP_1 \times 1.2 \\ 480 = CP_1 \times 1.2 \Rightarrow CP_1 = 400 \\ SP_2 = CP_2 \times 0.8 \Rightarrow CP_2 = 600 \\ 480 = CP_2 \times 0.8 \\ \hline TSP = 960 \\ \hline TCP = 1000 \end{array}$$

$$\text{Loss} = 40 \text{ Rs}$$

$$\text{Loss}\% = \frac{40}{1000} \times 100 = 4\%$$

MOHIT CHOUKSEY


0h

$$\text{Loss} = \frac{2 \times 20 \times 20 \times 480}{80 \times 120} = 40 \text{ ₹}$$

$$\text{Loss \%} = \frac{20 \times 20}{16} = 4\%$$


Pg 52
Q10

~~50%~~ ~~CP~~



hence $\frac{45}{150} \times 100 = 30\%$

T9



11.11 → 1/9

MP $\left(1 - \frac{10}{100}\right)$ ^{x.9}

MP $\left(1 - \left(\frac{1}{9}\right)\right)$ ^{11.11%}

= SP = CP $\times \left(1 + \frac{1}{7}\right)$ ^{14.28%}

MP $\times \frac{9}{9} = CP \times \frac{8}{7}$

MP = $\frac{9}{7}$ CP

MP = 1.2856 CP

MOHIT CHOUKSEY




$$80\% \xrightarrow{P} E \qquad 195 \xrightarrow{P} E \& S$$

$$70\% \xrightarrow{P} S$$

$$15\% \xrightarrow{F} E \& S$$

-24+21

SIR



$$n(A \cap B) = n(A) + n(B) - n(A \cup B)$$

$$\downarrow \qquad \downarrow \qquad \downarrow$$

$$85\% = 80\% + 70\% - x\%$$

$$x = 65\%$$

$$65\% \text{ of } T = 195$$

$$T = 300$$

RATIO

comparison b/w 2 quantities

Q.7 A student scored marks in 5 subjects in the ratio of 5:6:7:8:9. If the maxm. marks for all subjects is same and on aggregate, he scored 60% marks. in how many subjects did he pass the exam if passing marks is 50%.

Sol let the maxm. marks in each subject = 100
Total semester = 500

~~$$\text{He scored} = \frac{5}{100} \times \frac{6}{100} \times \frac{7}{100} \times \frac{8}{100} \times \frac{9}{100} = 300$$~~

~~$$5x + 6x + 7x + 8x + 9x = 300$$~~

$$35x = 300$$

$$x = 60/7$$

Annotations: $5 \times \frac{60}{7}$ and $6 \times \frac{60}{7}$ with arrows pointing to the first two terms of the crossed-out equation.

MOHIT CHOUKSEY

PROPORTION

I II III IV
 $a : b :: c : d$

$\frac{a}{b} = \frac{c}{d}$

$a \times d = b \times c$

if a, b, c, d are in continuous proportion.

$\frac{a}{b} = \frac{b}{c} \Rightarrow b^2 = ac \Rightarrow b = \sqrt{a \times c}$

b is GM (geometric mean)
 or MP (mean proportion) b/w (a & c)

DIRECT PROPORTION

$\uparrow a \propto b \uparrow$


$a = kb'$

$a/b = k$

division constant
 $\rightarrow a_1/b_1 = a_2/b_2$

\rightarrow unitary method

$\rightarrow y = mx$



ONGC 2012

Q Reduction in speed of a Railway engine is directly \propto to the sq. root of no. of compartments attached. If the maximum speed of the engine was 42 kmph when no compartment was attached and speed was 24 kmph when 9 compartments were attached. then the maxm. no. of compartments that can be carried forward by the Engine.

$42 - k(3) = 24$

Sol

~~$(x) \propto y^{1/2}$~~

Redn $\propto \sqrt{n}$

Redn = $k\sqrt{n}$

$Sp = Sp_{max} - k\sqrt{n}$

$Sp = 42 - k\sqrt{n}$

$24 = 42 - k\sqrt{n}$

$k=6$

$Sp = 42 - 6\sqrt{n}$

$6\sqrt{n} = 42$

$\sqrt{n} = 7$

$n = 49$

engine stops so at $n = 48$ engine max. speed.

Redn $\propto \sqrt{n}$

$$(v_2 - v_1) = k\sqrt{n}$$

$$42 - 24 = k\sqrt{9}$$

$$k = 6$$

$$42 - \cancel{0} = 6\sqrt{n}$$

$$n = 49$$

INVERSE PROPORTION

$\uparrow a \propto \frac{1}{b} \downarrow$

$a = \frac{k}{b}$

$a \times b = k$

$a_1 \times b_1 = a_2 \times b_2$

~~unitary method.~~

$x \times y = c$

never applicable

Rat. hyperbola

x-axis, y-axis asymptotes


if 40m \rightarrow 100 day
20m \rightarrow x = 200 //

CHAIN RULE

m	d	l	b	h	
30	15	40	60	90	$\frac{DP}{\frac{a}{b}} = k$
20	x	50	45	80	
					$\frac{IP}{a \times b} = k$
d \uparrow	m \downarrow	l \uparrow	b \uparrow	h \uparrow	
(15)	(30)	40	60	90	
(x)	(20)	50	45	80	

$$\frac{15 \times 30}{40 \times 60 \times 90} = \frac{x \times 20}{50 \times 45 \times 80}$$

MOHIT CHOUKSEY



1/8



P	Q	R	S
50	40	30	20

SIR

$$D \propto \begin{matrix} P \\ t \\ g \end{matrix} \quad D = k p g t$$

varies proportionately → graphs

↑ D ∝ angle

→ ↑ Growth
growth of a single microbe surviving human immunity system
within 24 hrs of entering the body

→ ↑ Potency (probability of microbe overcoming H. immunity man)

→ ↓ 1/E Toxicity (milligram of M/c req.)

$$D = \frac{p g}{t} k \quad \text{formulae}$$

LR DI
Logical Reasoning Data interpretation

dangerous level

$$D_p = \frac{5^2 \times 4}{8 \times 2} = 12.5$$

$$D_q = \frac{4^2 \times 5}{6} = 13.33$$

$$D_r = \frac{3^2 \times 4}{3} = 12$$

$$D_s = \frac{2^2 \times 8}{2} = 16$$

Pg no.
50
Q3

$$25p \rightarrow \frac{1}{5m} \text{ rs}$$
$$10p \rightarrow \frac{1}{10} \text{ rs} = 61$$

Q6

$$\frac{a}{b} = \frac{c}{d}$$

Q8

Profits A	=	$\frac{IA \times TA}{IB \times TB}$
Profit B		

investment → Time

Q7

$$\frac{a}{b} = \frac{b}{c} \quad \checkmark \quad \frac{36}{48} = \frac{48}{72}$$

Q9

$$0.7, 2.8$$
$$MP = \sqrt{0.7 \times 2.8}$$

$$\frac{In A}{In B} = \frac{5 \times 8 + 4 \times 4}{6 \times 4 + 3 \times 8}$$
$$\frac{5x}{8x}$$

$$\frac{8 \text{ months} \times}{T_B} \times \frac{5x}{8x} = \frac{1}{2}$$

$$T_B = 10 \text{ months}$$

Q

Prepp
Your Personal Exam Guide

MOHIT CHOUKSEY

12/8/16

SPEED, DISTANCE, TIME

$$S = \frac{D}{t}$$

$$\frac{1 \text{ km}}{\text{hr}} = \frac{1000 \text{ m}}{60 \text{ sec} \times 60 \text{ sec}}$$

$$= \frac{5}{18} \text{ m/sec}$$

(∵ $t = k$) $\uparrow S \propto D \uparrow$

$$\frac{S_1}{S_2} = \frac{D_1}{D_2}$$


(∵ $D = k$)

$\uparrow S \propto \frac{1}{t} \downarrow$

$$S \times t = k$$

$$S_1 \times t_1 = S_2 \times t_2$$

Q Gate 2017
2m



$$\frac{S_p}{S_c} = ?$$


~~$$S = \frac{D}{T}$$

$$C \rightarrow S = \frac{D_T}{T_1}$$

$$\frac{D_T}{T_1}$$


$$\frac{4/3}{T_1}$$~~

MOHIT CHOUKSEY



$$\frac{S_{PT}}{S_{PC}} = \frac{x}{L/3} = \frac{x+L}{2L/3} \Rightarrow 2x = x+L \Rightarrow x=L$$

$$\frac{L}{L/3} = \left(\frac{3}{1}\right)$$



$$\frac{S_{PT}}{S_{PC}} = \frac{L}{L/3} = \frac{3}{1} \quad (\because t = k)$$

they reach simultaneously (E=K=1/2)

AVERAGE SPEED

$$= \frac{\text{Total Distance}}{\text{Total time}}$$


AM > Gm > HM

$$\text{Avg sp} = \frac{TD}{TT} = \left[\frac{D_1 + D_2 + D_3}{t_1 + t_2 + t_3} \right]$$

$$\left[\frac{S_1 \times t_1 + S_2 \times t_2 + S_3 \times t_3}{t_1 + t_2 + t_3} \right] \checkmark$$

$$\frac{D_1 + D_2 + D_3}{\left(\frac{D_1}{S_1} + \frac{D_2}{S_2} + \frac{D_3}{S_3} \right)} \checkmark$$

$\times 50$
 $\times 52.5$
 $\times 51.5$
 $\times 48 \checkmark$



$$\text{Avg sp} = \frac{TD}{TT}$$


$$= \frac{2D}{D/a + D/b}$$

$$= \frac{2ab}{a+b}$$

harmonic mean of a & b.

Avg sp.


$$\text{Avg sp} = \frac{2ab}{a+b}$$



Average speed = $\frac{TD}{TT} \rightarrow \frac{\text{Area under any (s-t) graph}}{(TT)}$

Avsp = $\frac{50}{15} = 3.\bar{3} \text{ (m/s)}$ (during entire journey)

Q34
Pg 72



Avq sp = $\frac{1/2 + 1/2}{1/2 + \frac{x}{30} + \frac{1/2 - x}{20}}$

~~$\frac{1/2}{60} + \frac{x}{30} + \frac{1/2 - x}{20}$~~

~~$= \frac{1 + 4x + 6 - 6x}{120}$~~

~~$= \frac{1 + (-2x) + 6}{120}$~~

~~$\frac{7 - 2x}{120}$~~

~~$\frac{6}{120}$~~

(34)

120 kms


$\left(\frac{60}{60} + 30/30 + 30/10\right) \text{ hrs} = 24 \text{ kmph}$

(40)


$\frac{(8 + 6 + 16) \text{ km}}{(\frac{1}{4} + \frac{1}{4} + \frac{1}{4}) \text{ hrs}} = \frac{30 \text{ km}}{3/4 \text{ hrs}} = 40 \text{ kmph} \checkmark$

MOHIT CHOUKSEY

Q154




$\frac{1}{2} + 1 + \frac{1}{2} + 1 + \frac{1}{2} \times 2 + \frac{1}{2} + \frac{1}{2}$
 $1 + 2 + 1 + 1 = 5$ ✓




$\frac{\sqrt{100m}}{(15-10)m/sec} = 20sec$ Relative speed

127




$\frac{SL}{min}$
 f_{min}
 $D = \frac{S}{T}$
 $S_0 =$
 each leap = 8m (Tiger)

SIR



MOHIT CHOUKSEY

Q13
Pg 54




Bird remain in
here and do
not for
motion in
4 hrs

$$R_s = \frac{480 \text{ km}}{(70+50) \frac{\text{km}}{\text{hr}}} = \frac{480}{120} = 4 \text{ hrs}$$

$\frac{\text{km}}{\text{hr}} \cdot 100 \times 4 \text{ hrs} = \frac{400 \text{ km}}{\text{km}}$

DB = $SP_B \times t$
= 400 km



$$\frac{S_{PA}}{S_{PB}} = \sqrt{\frac{y}{x}}$$

① % ②

x & y are not
general time
taken.

Before meeting

x, y are time taken after meeting.

$$S_{PA} \times t = AC$$

$$S_{PB} \times t = BC$$

After meeting

A goes CB in 'x' hrs,

$$S_{PA} = \frac{CB}{x} = \frac{S_{PB} \times t}{x} \quad \text{--- ①}$$

B goes CA in 'y' hrs,

$$S_{PB} = \frac{AC}{y} = \frac{S_{PA} \times t}{y} \quad \text{--- ②}$$


T12 →

MOHIT CHOUKSEY

RACES. → pure application of Ratio / working but ratio

A beats B by 10m in a 100m race.

B = 90m → only significant meaning.
A = 100m






$$\left[\frac{S_A}{S_B} = \frac{100}{90} \right] (\because t = k)$$

Q A finishes 12m ahead of B and 18m ahead of C. while B finishes 8m ahead of C. then the length of the race.

$\frac{a}{36}, \frac{b}{48}, \frac{c}{60}, \frac{d}{72}$

Sol → Q2

Q1 Pg 53



$$\frac{S_A}{S_B} \times \frac{S_B}{S_C} = \frac{200}{18} \times \frac{25}{24} = \frac{125}{36}$$

$$\frac{S_B}{S_C} = \frac{250}{240} = \frac{25}{24}$$

$$\frac{125}{36} \times \frac{24}{25} = \frac{125}{36} \times \frac{4}{5} = \frac{125}{45} = \frac{25}{9}$$

$$\frac{S_A}{S_B} = \frac{200}{180} = \frac{20}{18}$$

MOHIT CHOUKSEY

Q1 Sol

Pg 53

$$\frac{A}{B} = \frac{200}{180} \quad \left[\frac{A}{C} = \frac{A}{B} \times \frac{B}{C} \right]$$

$$\frac{B}{C} = \frac{250}{240} \quad \frac{A}{C} = \frac{20}{18} \times \frac{25}{24}$$

$$\frac{A}{C} = \left(\frac{500}{432} \right) \begin{matrix} \times 2 = \\ \times 2 \end{matrix} = \frac{1000}{-864} = \frac{136 \text{ m}}{1000} \checkmark$$

Q2

$$\frac{A}{B} = \frac{L}{L-12}$$

$$\frac{A}{C} = \frac{L}{L-18}$$


$$\frac{B}{C} = \frac{L}{L-8}$$

$$\frac{L}{L-18} = \frac{L}{L-12} \times \frac{L}{L-8}$$

$$\frac{L}{L-18} = \frac{L}{L-12} \times \frac{L}{L-8}$$

$$\left[\frac{L-12}{L-18} = \frac{L}{L-8} \right]$$


use options $\rightarrow L=48$



$$\frac{B}{C} = \frac{36}{30} = \left(\frac{6}{5} \right) \times \frac{8}{8} = \frac{48}{-40} = \frac{8}{m}$$

MOHIT CHOUKSEY

CIRCULAR RACES



$$A = 15\text{m/s}$$

$$B = 10\text{m/s}$$

1) Time taken for meeting @ Start Point for the first time

$$\text{LCM}(t_{A1}, t_{B1})$$

$$\text{LCM}\left(\frac{\text{circumference}}{sp_A}, \frac{\text{circum}}{sp_B}\right)$$

$$\text{LCM}\left(\frac{30}{15} + \frac{30}{10}\right) = 6\text{sec}$$

@ 6 sec

$$DA = 15 \times 6 = 90\text{m} = 3\text{h}$$

$$DB = 10 \times 6 = 60\text{m} = 2\text{h}$$

@ 12 sec

$$DA = 15 \times 12 = 6\text{h}$$

$$DB = 4\text{h}$$

} @ SP

2) Time taken for meeting for the 1st time

$$\frac{\text{circumf.}}{\text{Rel.}(sp_A \pm sp_B)} = \frac{30}{(15+10)} = 1.2\text{secs}$$

@ 1.2 sec


$$DA = 15 \times 1.2 = 18\text{m}$$

$$DB = 10 \times 1.2 = 12\text{m}$$


@ 2.4 sec

$$DA = 36\text{m}$$

$$DB = 24\text{m}$$



Pg 53
 (7) x



A → 6 km/h
 B → 12 km/h

$$\frac{6 \text{ km}}{(6+12) \frac{\text{km}}{\text{h}}} = \frac{1}{3} \text{ h} \approx 20 \text{ min} \checkmark$$

$\frac{30}{18}$ ~~30~~

(8) @SP $\text{LCM}\left(\frac{600}{15}, \frac{600}{20}\right) \frac{\text{m}}{\text{sec}} = \frac{600}{5} = 120 \text{ sec} \approx 2 \text{ mins} \checkmark$

3rd formulae

No. of distant int meeting points on the track

$= \frac{P}{Q}$
 → value after ①
 → value after ② formulae
 (1.2) → 6

4th formulae

Time taken for meeting at the start point is independent of the dirn. of the runners.

5th formulae


if 3 Runners. $\text{LCM} \rightarrow (t_A, t_B, t_C)$

$$\text{LCM} \left[\frac{circ}{(A \pm B)}, \frac{circ}{(B \pm C)} \right]$$

Q Time taken \rightarrow train passes poll $\rightarrow \frac{L_T \leftarrow \text{Train}}{SP_T}$
 \rightarrow Platform $\rightarrow \frac{L_T + L_P \leftarrow \text{platform}}{SP_T}$
 \rightarrow to cross each 2 trains \rightarrow other $\rightarrow \frac{L_1 + L_2}{SP_1 \pm SP_2}$


MOHIT CHOUKSEY

10




$$\frac{L_2}{(s_1 + s_2)}$$

11



$$\frac{L_2}{s_2 - (s_1 + m)}$$

*



↓ sp = (b + c) → time less

↑ sp = (b - c) → time more

Pg 53
Q2



$$\frac{D}{\downarrow (20 + c)} = \frac{1}{3}$$

$$\frac{20 - c}{20 + c} = \frac{2}{3}$$

$$c = 4$$

$$\frac{D}{\uparrow (20 - c)} = \frac{1}{2}$$

MOHIT CHOUKSEY

Q74
Pg 77

$x = 8 \text{ km/h}$

$$x - y = \frac{D}{3t} \quad \left| \quad x + y = \frac{D}{t} \right. \quad \frac{\frac{D}{3t}}{\frac{D}{t}}$$

$$\cancel{x - y = \frac{D}{3t}} \quad \cancel{(x + y) = \frac{D}{t}}$$

$$3(x - y) = (x + y)$$

$$3x - 3y = x + y$$

$$2x - 3y = x + y$$

$$16 = 4y$$

$$y = 4$$

SIR

$$\frac{D}{\downarrow(8+c)} = t$$

$$\frac{D}{\downarrow(8-c)} = 3t$$

$$\frac{8-c}{8+c} = 3$$

Pg 84
Q9

use options
put $v = 60$

through
options

$$\frac{840}{v} - \frac{840}{v+10} = 2$$

$$\frac{840}{60} - \frac{84}{7} = 2$$

$$14 - 12 = 2 \quad \checkmark$$

MOHIT CHOUKSEY

CLOCK

Clock is an application of circular Race b/w hour hand and minute hand.

Min. hand

60 min \rightarrow 1 round $\rightarrow 360^\circ$
 1 min $\rightarrow \frac{1 \text{ round}}{360} \rightarrow (6^\circ)$

for RA of $(5\frac{1}{2})$ Min hand goes (6).
 for (RA of (1)) min hand $\rightarrow (\frac{12}{11})^\circ$

Hr. hand

12 hrs $\rightarrow 360^\circ$
 (60 min) \approx 1 hr $\rightarrow 30^\circ$
 1 min $\rightarrow (1/2)^\circ$

$5\frac{1}{2} \rightarrow 6$
 $\frac{11}{2} \rightarrow 6$
 $11 \rightarrow 12$
 $1 \rightarrow \frac{12}{11}$

Relative (RA) = $(5\frac{1}{2})^\circ$
 gain which the Min. hand over Hr. hand

Q. 1st variety \rightarrow 12 hrs

① coincide $\rightarrow 11$	} <u>24 hrs</u>	22] <u>day</u>
② right angle $\rightarrow 22$		44	
③ opposite $\rightarrow 11$		22	

FORMULAS

(x) & $(x+1)$ 0' clock

$5x \times \frac{12}{11}$ \leftarrow coincidence

$(5x \pm 15) \frac{12}{11}$ \leftarrow opposite \leftarrow Rt. angle

$(5x \pm 30) \frac{12}{11}$ \leftarrow ~~stand~~ opposite
 $x > 6$ (-)
 $x < 6$ (+)

$20 \times \frac{12}{11} = \frac{240}{11} = 21 \frac{9}{11} \rightarrow 4:21 \frac{9}{11}$ (win/loss)

$5x \rightarrow$

$5x \times \frac{12}{11} = \frac{60}{11} = 5 \frac{5}{11} \rightarrow 4:5 \frac{5}{11}$ (rt. Lnd)

$(5x \pm \frac{30}{2}) \times \frac{12}{11} = \frac{420}{11} = 38 \frac{2}{11}$

$(20+15) \times \frac{12}{11} = \frac{420}{11} = 38 \frac{2}{11}$

$(5x \pm 30) \times \frac{12}{11} = \frac{600}{11} = 54 \frac{6}{11} \Rightarrow 4:54 \frac{6}{11}$ (opposite)

Q if b/w 7 and 8

$(5 \times 7 - 30)$

$35 - 30 \quad 5 \times \frac{12}{11} = \frac{60}{11} = 5 \frac{5}{11} \checkmark$

$7:5 \frac{5}{11}$

- * $6^\circ \rightarrow 1 \text{ min}$
- $1^\circ \rightarrow (\frac{1}{6}) \text{ m}$

$\left[5x + \left(\frac{0^\circ}{6} \right) \text{ m} \right] \frac{12}{11} \leftarrow \text{Coin } 0^\circ$

$\left[5x \pm \left(\frac{90^\circ}{6} \right) \text{ m} \right] \frac{12}{11} \leftarrow \text{LL } 90^\circ$

$\left[5x \pm \left(\frac{180^\circ}{6} \right) \text{ m} \right] \frac{12}{11} \leftarrow \text{opp } 180^\circ$

$x \geq 6 (-)$
 $x \leq 6 (+)$

MOHIT CHOUKSEY

$$\left[5x \pm \left(\frac{D^\circ}{6} \right) \right] \times \frac{12}{11}$$

Pg 56
Q6

~~40°~~

~~$$\left[5x - \frac{D^\circ}{6} \right] \times \frac{12}{11}$$~~

~~5x~~

~~$$\left[5x2 + \frac{40}{6} \right]$$~~

$$\left[5x + \frac{D^\circ}{6} \right] \times \frac{12}{11}$$

$$\left[5x2 - \frac{40}{6} \right] \times \frac{12}{11} = \frac{40}{11} = 3\frac{7}{11} \Rightarrow 2 : 3\frac{7}{11}$$



or

$$\left[5x2 + \frac{40}{6} \right] \times \frac{12}{11} = \frac{200}{11} = 18\frac{2}{11} \Rightarrow 2 : 18\frac{2}{11}$$

Q What is the angle b/w the minute hand and hour hand at 9:25?

Sol


~~$$\left[5x + \frac{180}{6} \right] \frac{12}{11}$$~~
~~$$\left[5x - \frac{180}{6} \right] \frac{12}{11}$$~~
~~$$(125 - 30) \frac{12}{11}$$~~
~~$$95 \times \frac{12}{11}$$~~



$$\left[5x + \left(\frac{D^\circ}{6} \right) \right] \times \frac{12}{11} = 25^\circ$$


MOHIT CHOUKSEY

other



hr hand
 $1 \text{ min} \rightarrow \frac{1}{2}^\circ$
 $25 \text{ min} \rightarrow 12.50$

Mirror image \rightarrow symm. about y axis



Q: How much time / or minn hand hr hand kitni del baad mile?

$$60 \times \frac{12}{11} = \frac{720}{11} = \boxed{65 \frac{5}{11}} \text{ min letter (1 time)}$$

$$\sqrt{12 \text{ hr}} = 12 \times 60 = \frac{720 \text{ min}}{\frac{720}{11}} = 11 \text{ times}$$

MOHIT CHOUKSEY

1/14/2016

Aptitude and Reasoning

CAT - 30 to 35 Qn.

P and C (Permutation & Combination).

F.P.C. → Fundamental principle of counting

↳ 25 Qns out of 30 Qns

F.P.C. → Additive Rule → only one thing at a time

10 Boys 12 Girls

'a' monitor → 22 ways



$10 + 12 = 22 \text{ ways}$

'OR' → additive Rule can be applicable! hidden in the meaning of question.

Product Rule → 1B x 1G monitor

More than one thing
 $10 \times 12 = 120 \text{ ways}$

and
qn. → hidden or given or available



* Arrangement

${}^n P_r = \frac{n!}{(n-r)!}$

6 chairs, 6 members.

Ex:- $6 \times 5 \times 4 \times 3 \times 2 \times 1$

${}^6 P_6 = \frac{6!}{0!} = 720 \text{ ways}$ if 2 chairs broken

${}^6 P_4 = \frac{6!}{2!} = 360 \text{ ways}$

Q → {a, b, c} → {ab, bc, ca}

selen. ${}^n C_r = \frac{n!}{(n-r)! \times r!} \Rightarrow {}^3 C_2 = \frac{3!}{1! \times 2!} = 3 \text{ ways.}$

MOHIT CHOUKSEY

Q 12 people (handshake)

$${}^{12}C_2 = \frac{12!}{10! \times 2!} = \frac{12 \times 11}{2} = 66$$

$${}^n C_2 = \frac{n(n-1)}{2}$$

Q 12 points (str. line)

$${}^{12}C_2 = 66$$

* ${}^n C_r = {}^n C_{n-r} \rightarrow$ Ex:- ${}^5 C_2 = {}^5 C_3$
 ${}^8 C_5 = {}^8 C_3$

Q1> All 6 digit natural no.'s are being formed from 1st 6 natural no.'s without repetition. (w.r. R^n). How many such no.'s are divisible by 4?

Q2> How many 4 digit no. can be formed with 10 digits 0, 1, ..., 9. If no number can start with zero and if repetition are not allowed?

Q3> given digits 2, 2, 3, 3, 3, 4, 4, 4, 4. How many distinct 4 digit no.'s greater than 3000 can be formed?
 (Gate 2010) (a) 50 (b) 51 (c) 52 (d) 54.

Q4> All 4 digit natural no.'s are being formed from 1st five natural numbers. How many such no.'s are divisible by 4.

Me

Sol> ① $\frac{1}{\times} \frac{2}{\times} \frac{3}{\times} \frac{4}{\times} \frac{5}{\times} \frac{2}{\times} = 240$

1, 2, 3, 4, 5, 6

$$\begin{array}{r} 24 \\ \times 5 \\ \hline 120 \end{array}$$

② 0, 1, 2, 3, 4, 5, 6, 7, 8, 9

$\frac{9}{\times} \frac{9}{\times} \frac{8}{\times} \frac{7}{\times}$

$5^6 \times 4 \times 3 \times 12$

③ $\frac{2}{\times} \frac{7}{\times} \frac{6}{\times} \frac{5}{\times} = 5103 = 420$
 1, 2, 3, 4, 5


④ $\frac{5}{\times} \frac{5}{\times} \frac{5}{\times} \frac{1}{\times} = 250$

~~120~~

MOHIT CHOUKSEY

- Six → ① 192
 ② 4536
 ③ 51
 ④ 125

explanations




TU	TU	20	28
12	36	40	48
16	52	60	68
24	56	80	88
32	64	08	88
		04	8
		8	

44 ✓


⇒ $4 \times 3 \times 2 \times 1 \times 8 = 192 \text{ ways}$

② $\frac{9}{(1+9)} \times \frac{9}{(0+9)} \times \frac{8}{(0-9)} \times \frac{7}{(0-9)} = 4536$
 1, 2, 3, 4, 5



125 Ans

- ③ (2, 2) (3, 3, 3) 4, 4, 4, 4




① 3222 but two 2's are allowed → invalid no. also 3333 ✓

other way


$\frac{2}{3/4} \frac{3}{3} \frac{3}{2/3/4} \frac{3}{2/3/4} = \frac{54}{-3} = 51$


Q There are 12 towns equally to be divided into 4 zones. each town is connected to every other town in the same zone by 3 direct lines and each town is connected to every other town outside the zone by single direct line. How many lines are to be laid/built?

Sol 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12




MOHIT CHOUKSEY






Now



Now




Now



Q > 10 identical Balls are to be distributed among 3 friends. In how many ways can the distribution be done?

Sol > whole no. soln.



MOHIT CHOUKSEY

$$a + b + c = 10$$

$$\begin{array}{r} 1 \\ \hline 9 \quad 0 \end{array} \left. \begin{array}{l} 0 \\ 2 \\ 8 \end{array} \right\} \rightarrow 10$$

$$\begin{array}{r} 2 \\ \hline 8 \quad 0 \end{array} \left. \begin{array}{l} 0 \\ 8 \\ 7 \end{array} \right\} \rightarrow 9$$

$$\begin{array}{r} \hline 10 \quad 0 \quad 0 \end{array} \left. \begin{array}{l} 1 \\ 0 \\ 0 \end{array} \right\} \rightarrow 1$$

$$11 + 10 + 9 + 8 + 7 + 6 + 5 + 4 + 3 + 2 + 1 = 66$$

(or)

$$\frac{11 \times 12}{2} = 66 \checkmark$$

$$\frac{n(n+1)}{2}$$

Shortcut CONDITIONAL
soln.

N. No. soln

$$a + b + c = 10$$

$$\uparrow \quad \uparrow \quad \uparrow$$

whole No. soln. (\because 1 Balls all have)

$$A + B + C = 7$$

hence,

$$7 + 3 - 1 \quad C_{3-1}$$

$$9 C_2 = \frac{9 \times 8}{2} = 36 \checkmark \text{ Ans.}$$

0 Balls can be possibly assigned to anyone

Shortcut

① $n \rightarrow$ identical objects
② whole no. soln.
 $C_{(n-1)}$ applicable

whole no. soln. \rightarrow means can give 0 ball also.

$n \rightarrow$ identical objects
 $k \rightarrow$ no. of people.

here sol $(10 + 3 - 1) C_{(3-1)}$

$$= 12 C_2 = \frac{12 \times 11}{2} = 66.$$

Now

Natural No. soln.

$$a + b + c = 10$$

$$\begin{array}{r} 1 \\ \hline 8 \quad 1 \end{array} \left. \begin{array}{l} 1 \\ 2 \\ 7 \\ \vdots \\ 8 \end{array} \right\} \rightarrow 8$$

$$\begin{array}{r} 2 \\ \hline 7 \quad 1 \end{array} \left. \begin{array}{l} 1 \\ 2 \\ 6 \\ \vdots \\ 7 \end{array} \right\} \rightarrow 7$$

$$\frac{8 \times 9}{2} = 36$$

$$\begin{array}{r} 7 \\ \hline 2 \quad 2 \end{array} \left. \begin{array}{l} 1 \\ 2 \\ 1 \end{array} \right\} \rightarrow 2$$

$$\begin{array}{r} 8 \\ \hline 1 \quad 1 \end{array} \left. \begin{array}{l} 1 \\ 1 \end{array} \right\} \rightarrow 1$$

Noneed

$$\left\{ \because n C_2 = \frac{n(n-1)}{2} \right\}$$

MOHIT CHOUKSEY

Q 15 identical Balls are to be distributed among 4 friends (A, B, C, D) such that A should get atleast 3 balls, B atleast 2, C atleast 1. In how many ways can the distribution be done

Sol

$$\begin{matrix} A & + & B & + & C & + & D & = & 15 \\ 3 & \rightarrow & 2 & \rightarrow & 1 & \rightarrow & 0 & & \end{matrix}$$

$$15 - 6 = 9$$

$$A + B + C + D = 9$$

$$\begin{aligned} & {}^{9+4-1}C_{4-1} = {}^{12}C_3 = \frac{12!}{3!9!} = \frac{12 \times 11 \times 10 \times 9!}{9! \times 3!} \\ & = \frac{12 \times 11 \times 10}{6} = 220 \end{aligned}$$

MOHIT CHOUKSEY

* GEOMETRICAL P and C :-

12 points
~~00000~~

st. lines

$$\begin{aligned} & {}^{12}C_2 \text{ (if no points are collinear)} \\ & \text{---} \\ & {}^n C_2 \end{aligned}$$

$$\begin{aligned} & - {}^5 C_2 \text{ (5 points are collinear)} \\ & + 1 \text{ (one line possible)} \end{aligned}$$

12 points
~~00000~~

Δ 's



$$\begin{aligned} & {}^{12}C_3 \text{ (if no points are collinear)} \\ & \text{---} \\ & {}^n C_3 \end{aligned}$$

$$\begin{aligned} & - {}^5 C_3 \text{ (5 points collinearity)} \\ & \text{---} \\ & {}^r C_3 \end{aligned}$$

No. of diagonals of any 'n' sides polygon = $\frac{n(n-3)}{2}$

any 2 vertex makes hand shake
↓
side, Diagonal

$$\begin{aligned} & {}^n C_2 = \text{All sides} + \text{All Diag} \\ & \left\{ \begin{aligned} & {}^n C_2 - n = \text{All diagonals} \\ & \frac{n(n-1)}{2} - n = \frac{n(n-3)}{2} \end{aligned} \right. \end{aligned}$$




Q> If no. of diagonals of a n sided polygon is 50% more than its no. of sides then the polygon is —

Sol $\therefore 1.5x = \frac{x(n-3)}{2}$


$n-3 = 3$

$n=6$ → sides, 9 diagonals

S ↓
6
D ↓
9




Q> $211m$



how many Δ 's can we get from these 21 pts.

Sol $21C_3 - {}^{10}C_3 - {}^{11}C_3$
 $1330 - 1 - 1$

SIR



$\frac{{}^{11}C_2 \times 10 + {}^{10}C_2 \times 11}{550} = 445 = 1045 \checkmark$

$21C_3 - {}^{10}C_3 - {}^{11}C_3 = 1045 \checkmark$

{ all 3 should not from the same line }

MOHIT CHOUKSEY

Chess Board

$n+1C_2 \rightarrow n+1C_2 = \left[\frac{(n+1)n}{2} \right]^2$
 $\rightarrow 204 \text{ squares}$

$nC_2 = \frac{n(n-1)}{2}$

$9C_2 \times 9C_2 = 1296 \text{ Rectangles}$

1) Rectangles (n x n) $\rightarrow 9C_2 \times 9C_2 \rightarrow 1296$


$\sum n^3 = \left[\frac{n(n+1)}{2} \right]^2 = 1296$

2) Squares $\rightarrow 204$


$\sum n^2 = \frac{n(n+1)(2n+1)}{2} = 204$
 put $n=8$

3) different types of Rectangles

$\sum n = \frac{n(n+1)}{2} = 36$
 put $n=8$





area & perimeter same \rightarrow then same type



orientation differ.

$P=10$ $P=10$
 $A=4$ $A=10$

hence same Rectangle.




- $1 \times 1 \rightarrow 64 \rightarrow 8^2$
 - $2 \times 2 \rightarrow 7^2 \rightarrow 49$
 - $3 \times 3 \rightarrow 6^2 \rightarrow 36$
 - \vdots
 - $7 \times 7 \rightarrow 4^2 \rightarrow 22$
 - $8 \times 8 \rightarrow 1 \rightarrow 1^2$
- 204**

$1 \times 1, 1 \times 2, 1 \times 3, \dots, 1 \times 8 \rightarrow 8 \text{ types}$
 $2 \times 2, 2 \times 3, 2 \times 4, \dots, 2 \times 8 \rightarrow 7 \text{ types}$
 $3 \times 1, 3 \times 2, 3 \times 3, 3 \times 4, 3 \times 5, \dots, 3 \times 8 \rightarrow 6 \text{ types}$
 \vdots
 $8 \times 8 \rightarrow 1 \text{ type}$


all squares are included (जाने दो)

36


Gate 2017
2 marks



Redraw



CSAT (2011)



how many shortest routes are possible b/w A & B?

Sol

$$\frac{(R+C)!}{R! \times C!} = \frac{(5+3)!}{5! \times 3!} = 56.$$

Ex:- Apple = $\frac{5!}{2!} = 60$

BANANA = $\frac{6!}{3! \times 2!}$


[HHHHH VVV] = $\frac{8!}{5! \times 3!} = 56$

{HVNHNHNH}

MOHIT CHOUKSEY


* Linear Arrangement / Permutation :-

$3 \times 2 \times 1 = 3! = 6$ (circled)



circular Pn / circular arrangement:


$(n-1)!$
 $(3-1)! = 2!$ (circled) 2 ways



Q> A couple invited their 10 friends to a dinner party ^{to be held} across a circular dining table having 12 chairs such that there ~~has to~~ have to be exactly 1 friend b/w the couple.

Sol

$8!$ (with a scribble)




SIR



$2! (10 \times 9!)$

$2! \times 10!$
 $2 \times 10!$ (circled) Ans



H, W can interchange

Q> all 5 digit natural No.'s are being formed from 1st five natural no.'s without repetition. what is ^{sum} of all of those no.'s






$(n-1)! \times \underbrace{11111}_{n \text{ times}} (\sum d)^{\text{digits}}$
 $(5-1)! \times 11111 (1+2+3+4+5)$
 $4! \times 11111 \times 15$

CAT 2009
1, 3, 5, 7, 9

MOHIT CHOUKSEY

Q> vertex of a octagon are joined and Δ s are formed. How many Δ s are there whose vertex belongs to the vertex of octagon but none of its sides should belong to the side of octagon?

Sol SIR



me ✓

$$n=6$$

$$\Delta = \frac{n(n-5)}{3}$$

$$\frac{6 \times 1}{3} = 2$$

$$\frac{8 \times 2}{3} = \frac{16}{3}$$

$$\frac{8(3)}{3} = 8$$

$$\frac{5 \times 2}{3} = \frac{10}{3}$$


$$\frac{14}{11}$$

$8(3) \times$

$$\left. \begin{array}{l} n^2 - (n^2 - 2) \\ 36 - 34 \\ 2 \end{array} \right\}$$

$$\left. \begin{array}{l} 0 \downarrow 6 \\ 5 \downarrow 2 \end{array} \right\}$$

SIR



$$T(\Delta) \Rightarrow 8C_3$$

(no 3 collinear)

$$T(\Delta) = \Delta(1)$$


↓
common side

$$+ \Delta(2) (CDE)$$

$$+ \Delta(0) \rightarrow HCF$$

$$8C_3 = \Delta(1) + \Delta(2) + \frac{\Delta(0)}{?}$$

MOHIT CHOUKSEY




$$T(\Delta) = \Delta(1) + \Delta(2) + \left(\frac{\Delta(0)}{?}\right)$$

$$8C_3 = (4\Delta \times 8) + \binom{n}{8} + \Delta(0)$$

AB gone
HC also

AB side $\rightarrow 4\Delta$
8 side $\rightarrow 8 \times 4\Delta$



$$56 = 32 + 8 + \Delta(0)$$

$$\Delta(0) = 16$$

0 side common

$${}^nC_3 = n(n-4) + n + \Delta(0)$$

4 points gone

$$\Delta(0) = {}^nC_3 - n(n-4+1)$$

02/10/2016

PROBABILITY

Classical Defn:-

$$P = \frac{\text{favourable chances}}{\text{Total chances}} = \text{Probability}$$

Sample space = $\{1, 2, 3, 4, 5, 6\}$
in case of a dice

SS = $\{H, T\}$
in case of a coin

unbiased Events \rightarrow every event (equally likely)

- $P(1) = \frac{1}{6}$
- $P(2) = \frac{1}{6}$
- $P(3) = \frac{1}{6}$
- \vdots
- $P(6) = \frac{1}{6}$

- $P(H) = \frac{1}{2}$
- $P(T) = \frac{1}{2}$

Mutually exclusive events


and

Independent Events

{ Next Page }

MOHIT CHOUKSEY

✓ Mutually Exclusive events are events where happening of one event guarantees non-happening of the other.
 means $A \rightarrow$ happen, $B \rightarrow$ not happen.



for M.E.E.
Additive Rule $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 only one of the events happen @ time.

$P(A \cup B \cup C) = P(A) + P(B) + P(C)$

“oh”

English identifying key - word.

Q \rightarrow Dice

$P(\text{even}) + P(\text{odd}) - P(\text{even and odd}) = P(\text{even or odd})$
 $= 3/6 + 3/6 - 0 = P(\text{even or odd})$

$P(\text{or})$

✓ Independent Events are Events where more than one event can happen at a time without influencing the result of each other.

“and”

hint (1).

Ex:- Coin and dice is tossed simultaneously.

Product Rule $P(3m)$ and $P(t)$
 multiply \rightarrow tossed \rightarrow (tail)

$2/6 \times 1/2 = 1/6$

$P(A) \times P(B) \times P(C)$

Q $\rightarrow P(A) = 60\% \rightarrow$ A speaks Truth in 60% cases.

$P(B) = 75\%$

while answering the same qn. in either “Yes” or “No” they are likely to fight with each other in what % chances.

Sol: $P(A) = 3/5, P(\bar{A}) = 2/5$

$P(B) = 3/4, P(\bar{B}) = 1/4$

mutually exclusive

MOHIT CHOUKSEY

$$A \times \bar{B} + B \times \bar{A}$$

$$\frac{3}{5} \times \frac{1}{4} + \frac{3}{4} \times \frac{2}{5}$$

$$= \frac{9}{20} \approx \frac{45}{100} \approx 45\%$$

Q> There are 2 vacancies for which the husband and wife applied, $P(h) = 1/7 \rightarrow$ Probability of husband gets the job.
 $P(w) = 1/5$

only one gets the job	both	None	atleast one
?	?	?	?

Q> x is randomly chosen from 1st 100 natural no., what is the probability that chosen x satisfies the inequality

a) $\frac{28}{50}$

b) $\frac{29}{50}$

c) $\frac{59}{100}$

d) $30/50$

$$\frac{(x-40)(x-70)}{(x-30)} < 0$$

$$x \in [0, 100]$$

MOHIT CHOUKSEY

Q> A and B decided to meet b/w 6 and 7 p.m. on 14th Febr. 2017. what is the probability that they will meet provided one cannot wait for other for more than 20 minutes?

Q> Gate Qn.

Sol> ① $P(h) = 1/7$ $P(\bar{h}) = 6/7$
 $P(w) = 1/5$ $P(\bar{w}) = 4/5$

$$\left(\frac{1}{7} \times \frac{4}{5}\right) + \left(\frac{6}{7} \times \frac{1}{5}\right)$$

$$\frac{4}{35} + \frac{6}{35}$$

$$\frac{10}{35} \quad \text{⑩/35}$$

$$0.28$$

0.02
0.68
③ 1 hr 20 min 5/9
0.33×0.33
0.66
$0.25 \times \rightarrow 0.0625$

② $x-40 < 0$ $\frac{29}{50}$ ✓
 $x < 40, x < 70$
 $x < 30$

④ \rightarrow Gate $\frac{7}{16}$


SIR (1) $P(h) = 1/7, P(\bar{h}) = 6/7$
 $P(w) = 1/5, P(\bar{w}) = 4/5$

only one	both	none	@ least one
$h \times \bar{w} + \bar{h} \times w$	$h \times w$	$\bar{h} \times \bar{w}$	
$1/7 \times 4/5 + 1/5 \times 6/7$	$1/7 \times 1/5$	$6/7 \times 4/5$	
$\frac{10}{35}$	$\frac{1}{35}$	$\frac{24}{35}$	

$1 = \text{only one} + \text{both} + \text{None}$


$1 = \frac{10}{35} + \frac{1}{35} + \frac{24}{35}$

$1 - \frac{24}{35} = \frac{11}{35} \leftarrow P(\text{@least one})$



$\frac{(x-40)(x-70)}{(x-30)} < 0$

for $x > 70$



$x=40, x=70$ not allowed $\therefore < 0$

$\frac{\text{fav. chances}}{\text{Total chances}} = \frac{58}{100} = \frac{29}{50}$


41 to 69
29 values

signs can be putted on the no. line in alternate fashion.


(3) Time \rightarrow Real No.
 ∞ no. of values b/w 6 & 7.

$\frac{\text{favourable chances}}{\text{Total chances}} = \frac{\infty}{\infty} = \frac{f(A)}{T(A)}$
 favourable area / Total area

MOHIT CHOUKSEY



$$TA = 9$$



$$\text{area } (\Delta) = \frac{1}{2} \times 2 \times 2 = 2 \text{ units}$$

$$\text{unfavourable area} = 2 \times 2 = 4 \text{ units}$$

$$\text{favourable area} = 9 - 4 = 5 \text{ units}$$

$$\text{hence, } \frac{f(A)}{T(A)} = \frac{5}{9}$$

6:20 - 6:40 → Just a moment.

$$\text{here, } \Delta = \frac{1}{2} \times 40 \times 20 = 800 \text{ units}$$


$$\text{unfav Area} = 1600 \text{ units}$$

$$\text{Total area } TA = 60 \times 60 = 3600$$


$$\text{favourable area } fA = 3600 - (800 \times 2) = 2000$$

$$\frac{f(A)}{T(A)} = \frac{2000}{60 \times 60} = \frac{20}{36}$$

④



Red row



$$\frac{f(A)}{TA} = \frac{7}{16}$$

Conclusion

$$\text{if } TA \rightarrow \frac{3 \times 3 - 2 \times 2}{3 \times 3} = \frac{5}{9}$$

formulae
 $\frac{(2n-1)}{n^2}$

$$\text{if } TA \rightarrow \frac{4 \times 4 - 3 \times 3}{4 \times 4} = \frac{7}{16}$$

$$\frac{6 \times 6 - 5 \times 5}{6 \times 6} = \frac{11}{36}$$

MOHIT CHOUKSEY

Maths behind it

$$\left. \begin{aligned} 0 \leq x \leq 60 \\ 0 \leq y \leq 60 \end{aligned} \right\} TA$$

$$|x - y| \leq 20 \} f(A)$$



$$x - y = 20$$

if $x - y \leq 20$

if y comes first,

$$y - x = 20$$

$y - x \leq 20 \rightarrow$ To satisfy this ⁱⁿ equality, we have to move towards origin.



Do \rightarrow (33) (41) Pg 72

(33) 1 ----- 100

2 digit integers.

not divisible by 7

$$7 \times 7 = 49$$

~~7~~, 14, 21, 28, 35, 42, 49, 56, 63, 70, 77, 84, 91, ~~98~~

(14) no.

$$\frac{10}{7}$$

$$10 \text{ --- } 100$$

$$\begin{array}{r} \textcircled{91} \\ - 14 \\ \hline 77 \end{array}$$

SIR [1-100] \div by 7

$$\frac{100}{7} = 14 \quad [7, 14, 21, \dots, 98]$$

[10-99] \rightarrow Total No.'s = (90)

$$\text{div. by } 7 = (14 - 1) = (13)$$

$$\frac{f_c}{T_c} = \frac{77}{90} \checkmark$$

MOHIT CHOUKSEY

$11Y = 366d = 52 \times 7 + 2 \text{ odd day}$
 53rd saturday → 2 chances of saturday
 $\frac{fc}{TC} = \frac{2}{7}$ ✓
 P.S.
 S.S.
 S.M.
 M.T.
 T.W.
 W.T.
 T.F.


$\frac{2}{7}$ ← Ans.

* Sample space (dice) = {1, 2, 3, ..., 6}
 $P(\text{even}) = 3/6 \rightarrow \{2, 4, 6\}$
 $P(\text{prime}) = 3/6 \rightarrow \{2, 3, 5\}$

Q → (Conditional Probability Based).
 A dice is thrown at random. what is the probability of getting a prime no. on the dice provided the dice had shown an even number.

Sol → S.S._{new} = [2, 4, 6]
 $P(\text{prime}) = P\left(\frac{\text{prime}}{\text{even}}\right) = \frac{1}{3}$ ✓

Pg 71 (20) X, Y
 X → 60% → 96% reliable
 Y → 40% → 72% reliable



MOHIT CHOUKSEY

53

$\frac{10}{100} \rightarrow \text{HIV}^+$

HIV+ \rightarrow 95% (True)

HIV- \rightarrow 89% (-ve)

SIR

$$\frac{0.1 \times 0.95}{(0.1 \times 0.95) + (0.9 \times 0.11)}$$

\uparrow +ve \uparrow +ve \uparrow -ve \uparrow m/c \rightarrow +ve (-vector)
 (Note: The denominator terms are 0.1×0.95 and 0.9×0.11)

0.4896 Ans

Pq 54

- (5) P(2) = 1
- P(3) = 2
- P(4) = 3
- P(5) = 4
- P(6) = 5
- P(7) = 6
- P(8) = 5
- P(9) = 4
- P(10) = 3
- P(11) = 2
- P(12) = 1

- { 1, 6 }
- { 2, 5 }
- { 3, 4 }
- { 6, 1 }
- { 5, 2 }
- { 4, 3 }

Pq 54

(6) $\frac{1}{3} \times \frac{1}{4} \times \frac{1}{5} \times \frac{1}{6}$

independent events

SIR

None of them solves qn. \Rightarrow $\bar{A} \times \bar{B} \times \bar{C} \times \bar{D}$

Qn is not solved

Qn. is solved $\Rightarrow 1 - (\bar{A} \times \bar{B} \times \bar{C} \times \bar{D})$

(or) atleast one of them solves Question

All of them solves the qn. $\Rightarrow A \times B \times C \times D$

Q7

4 times
2H, 2T

- | | | | |
|------|------|------|------|
| HHHT | HHTH | HTHH | THHH |
| HHTT | HTHT | HTTH | THTH |
| HHTT | HTHT | HTTH | THTH |
| HTTT | HTTT | TTHT | TTTH |
| TTTT | TTTT | | |
- $\frac{4}{16} = \frac{1}{4}$

H	H	T	T
H	T	H	T
H	T	T	H
T	H	H	T
T	H	T	H
T	T	H	H


$\frac{6}{16}$ alter.

6C_2

alter. method $\rightarrow {}^6C_2 \left(\frac{1}{2}\right)^2 {}^2C_2 \left(\frac{1}{2}\right)^2$

$\frac{6}{16}$

Q 10 penalty shootouts. ${}^{10}C_4 \rightarrow$ chances in which goal happens.



${}^{10}C_4 \cdot (s)^4 \cdot (f)^6$
 \rightarrow success

Pg 90
 (177)

${}^{10}C_6 \times$

0.2508

Pg 54

(10) $\frac{0.04}{100} = \left(\frac{1}{25}\right)$

Pg 54

(1) $nC_2 = n \cdot \text{shakes}$

$\frac{n(n-1)}{2} = 153$

(18) ✓

(2) ✓



* **LOGICAL REASONING** *



4 Rules \rightarrow Rule 1 \rightarrow draw all possibilities/(Cases).
 Rule 2 \rightarrow for a statement to be True, it have to be true in all the cases.

Rule 3 \rightarrow If a statement is false even in one of the case, then it will be considered false forever.

Rule 4 \rightarrow Try to proof a statement false as early as possible.





Rules/General \rightarrow Read dirn's carefully \uparrow






no informn. about
Hens and Birds
hence
3 possibilities.

conc i ✗
ii ✓ (b)





i. ✓
ii ✗
(a) ✓




(i) ✗
(ii) ✗


(d) ✓




MOHIT CHOUKSEY




(i) ✗
(ii) ✓ (b) ✓





(i) ✓
(ii) ✓ (c)




SIR
(8)



i ✓
ii ✓ (d) ✓




(i) ✓
(ii) ✓ (d) ✓







(1) ✗
(2) ✗ (c) ✓



MOHIT CHOUKSEY




- (1) ✓
- (2) ✓
- (d) ✓







- (i) ✗
- (ii) ✗
- (iii) ✗
- (iv) ✗
- (d) ✓



- (i) ✗
- (ii) ✗
- (iii) ✓
- (iv) ✓
- (c) ✓




- (i) ✗
- (ii) ✓
- (iii) ✗
- (iv) ✓



MOHIT CHOUKSEY



136

SIR



No informⁿ about Manager and Executive

hence Cases





(i) ✗ (Bullshit)

(ii) ✗ —



(c) ✓

Pg no. 57

8 to 9




∴ all are adjacent



(11)


1, 2, 3, 4, 5, 6



1 → 2 ✗ 3 ✗ 4 ✗ 6 ✗
opp.
1 ← 5
opp. (a) ✓

MOHIT CHOUKSEY


* **DIRECTION** :-



Horizontal ^(H) = E⁺, W⁻
 Vertical ^(V) = N⁺, S⁻
 Apply pythagoras.


$$D = \sqrt{H^2 + V^2}$$

Q) Person goes 40m North take a Right turn goes 60m takes a left Turn ^{& goes} 50m and takes another " " " 45m



$$H = |60 + 45 - 95| = 10$$

$$V = |40 + 50 - 70| = 20$$






$$D = \sqrt{70^2 + 20^2}$$

$$D = \sqrt{500}$$

$$D = 10\sqrt{5} \text{ Ans} \therefore$$

MOHIT CHOUKSEY



Isosceles Right \triangle



$$\sqrt{a^2 + ?^2} = (2a)^2$$

$$a^2 + ? = 2a$$

\Downarrow



$$\sin 30^\circ = \frac{a}{h}$$

$$\frac{1}{2} = \frac{a}{h}$$

$$h = 2a$$


Pg 74

Q 52


$$+\sqrt{2} + 1 - \cancel{1} + \cancel{1}$$

$$+1 - \sqrt{2} + 1 - \cancel{1} + \cancel{1}$$

2



SIR



(149) \rightarrow H.W. ©

MOHIT CHOUKSEY

17/11/2016

Data Interpretation

$$\% \text{ change} = \left[\frac{\text{change}}{IV} \right] \times 100$$

$$\% \text{ change} = \left[\frac{FV - IV}{IV} \right] \times 100$$

FV → final value

IV → initial value

if % change → +ve → FV > IV → % ↑ → growth rate

→ -ve → FV < IV → % ↓ → decline rate

Ex (a) 40 → 20 ⇒ $\frac{-20}{40} = \left(-\frac{1}{2} \right) \times 100 \approx -50\% \downarrow$

(b) 40 → 50 → $\frac{10}{40} = \frac{1}{4} \approx 25\% \uparrow$

(c) 40 → 55 → $\frac{15}{40} = \frac{3}{8} \approx 37.5\% \uparrow$

Ans ∴
(a) in

% change maxm. → | change |

% ↑ is maxm. → +ve value in account.

% ↓ is maxm. → -ve value in account.

Ex :-

2015	2016	2017
50	60	72

$-\frac{10}{50} = -\frac{1}{5} \approx 20\% \uparrow \checkmark$

$50 \times 1.2 = 60 \times 1.2 = 72$

50 40 12

$50 \times 0.8 = 40 \times 0.8 = 32 \checkmark$

MOHIT CHOUKSEY

MOHIT CHOUKSEY

Rs 10,000

$\frac{90^\circ}{360^\circ} = \frac{1}{4} \rightarrow 25\%$

$\frac{1}{4}$ of 10,000 = 2500

1,00,00,000

100% \rightarrow 360°

1% \rightarrow $\frac{18}{360}$ or 3.6%

360° \rightarrow 100%

1° \rightarrow 5/18%

Pg 58 CH # 13

Q2 $\left[\frac{63^\circ - 36^\circ}{360} \right] = \frac{27}{36} = \frac{3}{4} \approx 75\%$

Q3

$\frac{81^\circ + 63^\circ}{360^\circ} = \frac{144^\circ}{360^\circ} = \frac{2}{5}$ of Total

$= \frac{2}{5}$ of (200)

$= 80$ Lakh

Method

40% of the Total \leftarrow other method

40% of T

Q6 0.04 of TCP = 15730

TCP = 893250

of 5500

Total selling Price (TSP) = $\frac{893250 \times 1.3}{5500} = 92.95$

loop

1% \approx 4 Thousand

4% TCP \approx 16 Th

TCP = 4 lakh

$$TSP = 4 \times 1.3 = \frac{5.242}{5.500}$$

lwpq

9

(10) ✓



(11) ✓

(12) ✓

(15) Total quantity = 5 lakh tonnes
Total Revenues = 250 Crore

Ratio of Revenue $\frac{1/Kg}{4/Kg}$

(15) SIR



$$I_1 = \frac{2 \times \frac{12\% \text{ of } R}{14\% \text{ of } E}}{2} = 4/1$$

$$I_4 = \frac{1 \times \frac{8\% \text{ of } R}{22\% \text{ of } E}}{2} = 4/1$$

MOHIT CHOUKSEY

Pg 77

Q 75

200 units

SIR

15 of TCP = 4.5 lakh

✓ TCP = 30 lakh
2012

✓ Profit 2012 = 10 lakh

T.S.P. 2012 = 40 lakh

$$S.P./\text{per unit} = \frac{40 \times 10^5}{200} = 20,000$$

Pg no. 60

Q 18

$$\frac{M_{2008}}{F_{2008}} = 2.5 \checkmark$$

assume $F_{2008} = 100 \checkmark$
and $M_{2008} = 250 \checkmark$ (Bec. ratio is fixed)

$$\frac{M_{2009}}{F_{2009}} = \frac{600}{200} \quad (3) \checkmark$$

(Bec. Ratio is fixed)

Final value of male $\frac{600 - 250}{250}$ initial value of male

$$= \frac{35}{25} = \frac{7}{5} = 1.4 \times 100 = 140\%$$

(19) 2012-2013 \rightarrow GDP \uparrow 7%
2012-2013 \rightarrow 50 to 60 USD

SIR	GDP in ₹	1 US \$	GDP in 1 US \$
Q19	GDP before 2012-2013	100 ₹	$\frac{100}{50} = 2 \$$
		50 ₹	
	GDP after 2012-2013	107 ₹	$\frac{107}{60} = 1.783 \$$
		60 ₹	

$$\left[\frac{2 - 1.783}{2} \right] \times 100 = -10.83\% \downarrow$$

20 ↑ → 16.8% ↓
 ↳ production purchase ↓. power

Q20

Type III	Type II	Type I	Type X
$\frac{46}{114}$	$\frac{40}{144}$	$\frac{40}{75}$	$\frac{40}{108}$

Pg 84
Q123

	(cm)	(kwh)
M	20 → 12	
N	45 → 25	
O	75 → 45	
P	100 → 57	

20 km → 12 kwh 45 → 25 75 → 45
 1 km = $\frac{12}{20}$ 1 km = $\frac{25}{45}$ 1 km = $\frac{45}{75}$
 0.6 0.5 0.6
 1 → $\frac{57}{100}$
 0.57

MOHIT CHOUKSEY

<u>SIR</u>		(km)	(kwh)
M	20	←	12
N	25	←	13
O	30	←	20
P	25	←	12

$$\left(\frac{13}{36}\right) \quad \left(\frac{13}{25}\right) \uparrow$$

(116) ✓ (87) ✓
Same

Pg 78 Q87 →

21x2	42	} α.
15x3	45	
23x2	46	
<u>42 + 45 + 46</u>		(15)
3		

SIR

21x2 = 42
15x3 = 45
23x2 = 46
<u>133</u>

(C)
(88)

Students Corrected
Class + W + NA = students
80 + 5 + 15 = 100
10 + 70 + 20 = 100

Q. ESE 2017

apna Qn	M	F	T
5L-	1(38, 38)	5(34, 40)	6
5-10L	1(32, 32)	8(35, 53)	9
10-15L	8(21, 65)	3(37, 13)	11
15+	2(32, 33)	2(27, 40)	4
Total	12	18	(30)

30 is needed

(i) The percentage of the people older than 35 years can be almost

Sol 8

Extend the Table

	M	F	T
	1	4	5
	0	7	7
	7	3	10
	0	1	1
			(23)

23/30 × 100 = 76.6%

(ii) Then the ^{minm} % of people age less than 40 years





M	F	T
1	1	2
1	1	2
1	1	2
2	1	3

MOHIT CHOUKSEY

Q
 10 samples P Q Y
 (70) | (57) | 82 | 84 | 98 | (66) | (34) | 87 | (79) | (71)

This shows the % of milk in each sample. If any two samples are mixed & form a new sample then on maxm., how many distant pairs of samples will never give a composition of more than 80% milk.

Sol



$\frac{{}^6C_2}{2}$
 $\frac{6!}{2! \cdot 4!}$
 SRS 15

Q — .

Prepp
 Your Personal Exam Guide

MOHIT CHOUKSEY


SI/CI

Simple Interest / Compound Interest

$$SI = \frac{P \times R \times T}{100}$$

$$y = mx$$

RS 100 @ 10%




(A)

$$\text{Amount} = P + SI$$

$$y = c + mx$$

RS 100 @ 10%



₹8P → 96yrs ↑

@ CI $P_1 + I_1 = A_1 = P_2$

$$P + \frac{PR}{100} = P \left(1 + \frac{R}{100}\right)^1 = A_1 = P_2$$

$$A_n = P \left(1 + \frac{R}{100}\right)^n$$

$$CI = A_n - P$$

Amount is compounded (half yearly)

$P, R = 10\%$ per year; $t = 2$ year

$$A_2 = P \left(1 + \frac{5}{100}\right)^4$$

(Amount is compounded Quarterly) @ +1yr

$P, R = 5\%$ per half yearly (phy), $t = 2y$

$$A_2 = P \left(1 + \frac{2.5}{100}\right)^8$$

$$* (CI - SI)_{2y} = P \left(\frac{R}{100}\right)^2$$

$$* (CI - SI)_{3y} = P \left(\frac{R}{100}\right)^3 + 3P \left(\frac{R}{100}\right)^2$$

Rs 100 @ 10%

	CI	SI
P	100	100
I_1	10	10
P_2	(100 + 10)	100
I_2	(10 + 10)	10

MOHIT CHOUKSEY

Q certain sum of money becomes 25 times in 48 yrs at a S.I. In how many yrs will it become 49 times at S.I. ?

Sol

$$\text{Amount} = P + SI$$

$$25P = P + 24P$$

$$25P = P + 24P$$

$$\sqrt{24P} \propto 48y$$

↑ SI ∝ t ↑

$$A = P + SI$$

$$49P = P + 48P$$

$$\left(\frac{PR}{100}\right) \times \frac{R}{100} = P \left(\frac{R}{100}\right)^2$$

Q Certain sum of money doubles itself in 5 yrs at C.I. In how many years will it become 8 times at C.I.

Q CI

'm' times in 'y' years
 $(m^n) \rightarrow (n \times y) \text{ years}$

 2 times in 5 years

 $8 \approx 2^3 \text{ (times) in } 3 \times 5 = 15 \text{ yrs}$

$$A = P \left(1 + \frac{R}{100}\right)^n \quad \rightarrow \quad 8P = P \left(1 + \frac{R}{100}\right)^{15}$$

$$2P = P \left(1 + \frac{R}{100}\right)^5$$

Cubing

$$8P = P \left(1 + \frac{R}{100}\right)^{15}$$

Q49
 Q79
 then
 pg 52
 Q2
 Q10

Q49 → ~~Smritition~~ → ~~20% annually~~

~~$A = (1.2)^n$~~
 ~~$f(n=3)$~~

$$A = P \left(1 + \frac{R}{100}\right)^n$$

$$P \left(1 + \frac{20}{100}\right)^n$$

$$A = (1.2)^n$$

$$f(n=3) = 1.728 P$$

$$f(n=4) = 2.07 P$$

$$A = P \left(1 + \frac{R}{100} \right)^n$$

$$2P = P \left(1 + \frac{R}{100} \right)^{10}$$

$$2^{1/10} = \left(1 + \frac{R}{100} \right)$$

Q2

$$A = P + SI$$

$$3080 = P + \frac{PR \times 3}{100}$$

$$3400 = P + \frac{PR \times 5}{100}$$

$$2600 + \frac{\text{interest}}{(160 \times 3)} = 3080$$

$$2600 + 800 = 3400$$

$$320 \leftarrow 2y(SI)$$

$$160 \leftarrow 1y(SI)$$

$$\frac{T_{10}}{5324} = P \left(1 + \frac{R}{100} \right)^3$$

$$4840 = P \left(1 + \frac{R}{100} \right)^2$$

Q A large cube was dipped in paint, taken out and then divided into 64 equal smaller cubes. how many cubes are painted on 3 sides, 2 sides, 1 side, 0 side.

Solution

$$T = (4 \times 4) \times 4 = 64$$


$$3S = 8$$

$$3 \text{ side} +$$

$$2S = 2 \times 12 = 24$$

$$1S = 4 \times 6 = 24$$

$$\frac{2S}{\text{side}} = 2 \times 12 = 24$$





$$+ OS = \text{Remaining} = 8$$

MOHIT CHOUKSEY

Q. A large cube was dipped in paint, taken out and then its length was divided into 5, width was divided into 4, height → into 3 equal parts. then, how many cuboids are painted on 3s, 2s, 1s, 0s.

Sol.



$$T = (5 \times 4) \times 3 = 60$$

$$3S = 8 \cdot +$$

$$2S = 4[(3) + (2) + (1)] = 24$$

$$1S = 2[(3) + (2) + 6] = 22$$

$$0S = 6$$

Q159 (56) ✓

$$T.S.A. = 6(\text{side})^2$$


$$6(4)^2$$

$$TSA = 96$$

Q121

MOHIT CHOUKSEY

(121)



$$6(\text{side})^2 = \text{TSA}$$

$$6(1)^2 \times 27 = \text{T.S.A.}$$

$$6(3)^2 = \text{V.A.}$$

$$54 \times 2 = 108 = \text{No. V.A.}$$

+, -, ×, ÷

↑ ↓ × ÷

Rule on Averages → ① If each and every $\frac{\text{set}}{\text{pr.}}$ is ↑, ↓, ×, ÷ by a constant, then their arithmetic mean is also ↑, ↓, ×, ÷ by the same constant.

② Sum of the deviations taken from arithmetic mean is equal to zero.

(19) Pg 71

$$\text{Standard deviation} = \sqrt{\frac{d_1^2 + d_2^2 + \dots + d_n^2}{n}}$$

* $d_1 = -2$ $d_2 = 0$ $d_3 = 2$
 * 1 3 5
 * 8 10 12
 $d_1 = -2$ $d_2 = 0$ $d_3 = +2$

$$\bar{x} = 3$$

$$\bar{x} = 10$$

Q47

a ✓
 b × avg atleast.
 d × avg every.

(b) ✓

* (T)

95% ile → 94.3, 94, 95
 Avg 93
 5% ile → 92.3, 92, 91

consistently high ↑

* (Q)

95% ile → 5.3, 5, 6
 Avg 4
 5% ile → 2, 3, 3.3

β	95%, 97, 96	highly inconsistent
95/ile	Avg 92	
5/ile	2, 1, 3	

Date 2016

(181)

37 (73)

2012 → M - W → 41°C
 T - T → 43°C
 T → 15% > M

15% of 41

SIR

$$\frac{M + T + W}{3} = 41$$

$$\frac{T + W + Th}{3} = 43$$

$$M + T + W = 123$$

$$T + W + Th = 129$$

$$Th - M = 6$$

$$Th = 1.15M$$

2 The average weight of 25 students was 42 kg's. Two new student having weight 54 and 66 kg joins the class. What's the new average.

Sol


$$\frac{\text{Sum}}{N} = \bar{x} \leftarrow \text{average}$$

$$\text{Sum} = N\bar{x}$$

$$\text{Sum} = 42 \times 25$$

$$= 1050 + 54 + 66$$



07



68	-2
75	+5
77	+3
72	-8
69	-1
74	+4

70 + 1/6
70.66


* Seating Arrangement





c is 2 places right of m.
r is — " — " left — " — "

- ① equal parts.
- ② @ centre.
- ③ R → Immediate Right.
L → Immediate left.



- ④ $\begin{bmatrix} ACB \\ BCA \end{bmatrix}$
- ⑤ (m n)
(n m)
- ⑥ $\leftarrow \begin{matrix} m \\ n \\ p \end{matrix}$



- T @ L
- S P @
- Q W @ L




9





br
ar

SIR



[GDB]
[BDE]




* BLOOD RELATIONS

There are 5 Rules :-

- 1) Draw family Hierarchy Tree
- 2) Keep on marking genders
- 3) Relationship
- 4) A+ wife M-
- 5) don't Assume don't Names


here gender is not known

A is the one level up in the family hierarchy



Either Nephew or Niece (D with S-)
or
C.B.D
cannot be determined.
but Nephew
Niece


Q1




Q1 ✓

Q2


Q1 Sir




women related to man



Q2 Sir




Q9



MOHIT CHOUKSEY

Analytical Reasoning

Pg 66



Ds support → Finance

E, F → marketing

F → operations ← C & E
support support

A → Finance & IT
main

③ cv ✓

④ operat @ ✓ ✗

⑤ A & @ ✓ ✗


categories here are only 2

SIR


co-ordinat support

A	Finance	IT
B	Finance	Marketing
C	Finance	opr
D	IT	Fin
E	mkty	opr
F	opr	mkty


[3 people - Co-ordinate - Fin]




⑨ ⑩ M, K, R



⑩ X, M, T



⑪ P, X, Q, S



categories 6

Pg no 66

	Hock.	volly	chick	base	foot
Q1 R	✓	✓		✓	
K	✓	✓	✓		
S	✓			✓	✓
Q		✓	✓	✓	✓
M				✓	✓

③ cv

④ shg

MOHIT CHOUKSEY

please write in short

- 1) Solio.
- 2) Educ.
- 3) Acc.
- 4) < English
- 5) Eco
- 6) Psycho
- 7) Hindi

Eco
Psycho
Hindi

5 Girls

K → 2nd height

Ascending

~~A → K < R~~
~~W → P > R~~
~~N → P > M~~
~~A → R & M~~
~~B → R > M~~
~~W → N > P~~
~~A → R > N~~

P > R
 P > M
 R > M
 N > P

~~N > P > R > M~~
 (d) 7 ✓
 K < R
 R & M = R = M
 R > N
~~K < R = M > N~~

C ✓

Rules (एगान दे कृपया) :-

B follow A
 OR
 A is followed by B

\downarrow
 BAM → A
 B:ISAM → B

A > B shortest
 B > C
 C > F

WB Pg 67 5 girls

~~R > P~~
~~K~~
 P > R
 P > M
 N > P

SIR solutions

Height
 1 2
 K
~~P > R~~
~~P > M~~
~~R > M~~
~~N > P~~

shortest one
 have to
 come from
 R and P

~~N > K > P > M > M~~
 asc ↑

जो पक्ष बाकी
 left side aa gaya
 तो count wala
 hoga
 shortest


Age K ≠ Youngest

~~K < R~~
~~P < M~~
~~R = M~~
~~R < N~~

$> P$
 $N > R = M > K > P$

1 2 3 4
 3 2 4
 ↓
 des

youngest
 one
 has to
 come
 from
 K
 P
 R



S
P
T
W

20 to 22

A K S, R N

R Y W G R B

R P O S W

K → S → Y

S → R → R* & W*

N → P → B & Y

A → W

R → Y or G

d ✓

	Red	Yellow	Blue	white	Green	
Amar					Red	Reading
Kap						Playing
Sas						outing
Roh						Singing
Nag						working

21

	Red	Yellow	Blue	white	Green	Read	Play	outing	Sing	working
Amar		X			X					✓
Kapil	—	X	—	—					✓	
Salvesh	✓					X				X
Rohan										
Nagesh	✓	X	X	—			✓			

N.P.

MOHIT CHOUKSEY

SIR's Solution

	colour	hobby
A	Yellow yellow	write ✓
K	Yellow	Sing ✓
S	Red ✓	Reads ✓ writes ✓
R	Yellow ✓ green ✓	Blue or white Read ✓
N	Blue ✓ Yellow ✓	white or green Play ✓


Not in gate

(23) R → Blue

(24) Kapil → can't be determined ✓

A → even
B → odd

Q44 → (Gate II)



Eswal does not live on floor number Bhola.

Don't tabulate.

(b) ✓ Bullshit

(50) 4 children

SOM < Riaz
Shiv < Ansu
Ansu < group
↓
youngest

SIR
SOM < Riaz ✓
Anshu < Shiv
shiv (or) Riaz

(a) ✓

	P	Q	R	S	T	U
P						
Q						
R						
S				✓		
T						
U						

don't Tabulate the data



- (a) ✗ R - Defense
- (d) ✗ R - Telecom
- (c) S & U can't be together

(b) ← Ans

Q161 Pg 88

$A > B$
 $B > C$
 $A > B > C$

Since data is quantitative




MOHIT CHOUKSEY

161 / sir (a) ✓

copy cohen
↑
Book ✓

Lettering

(1) A (2) B (3) C D - - E (5) - - F - - G - - H I J (16) K
 L M N (13) P Q R S T (20) U W
 V X Y (25) Z (26)



Pg no. 64

- Q5
- Q8
- Q12
- Q16



(5) 1 2 5
 AB = E
 EA = Y
 5 1 25

3 4 25
 CD = Y
 BC = M
 13

(8) CAT =

MOHIT CHOUKSEY

LETTERING



(15) ABK : V :: BCD : _____ 8
10

(12) ~~C~~ A ~~R~~ P ~~E~~ T : ~~T~~ C ~~E~~ A P R :: _____ : ~~L~~ N ~~A~~ A N ~~T~~ O ~~I~~

$\downarrow \downarrow$ $\downarrow \downarrow$
 5 20 20 5

(5) →

SIR (5)

A	B	=	E
1^2	$+ 2^2$		5

C	D	=	y
3^2	$+ 4^2$		= 25

E	A	≠	y
5^2	$+ 1^2$		= 26 = z

(12) ~~C~~ A ~~R~~ P ~~E~~ T : ~~T~~ C ~~E~~ A P R

: ~~T~~ C ~~E~~ A P R

~~N~~ A ~~T~~ I O ~~N~~ A L : ~~L~~ N ~~A~~ A N ~~T~~ O ~~I~~

A B K : 18
: 22

1 x 2 x 11

B C D

2 x 3 x 1 = 2X

B ← D → E
O ← R → P → E

(21) BOARD : CPBSE

CHAIR :

BHOPAL EERMDI

NAGPUR

Q21
22
24
25

SIR (BHPAL) : (EERMDI)

(2)

+3/-3/+3/-3/+3/

(22) MONKEY ← Y : X D J M N L

Tiger : 0

Y E K N O M
MONKEY : X D J M N L


MOHIT CHOUKSEY

10
← Best of luck : Kaly TSB
← Good wishes : shw DA
10 6

NS
Alg
Gate Qns
Doubt

Prepp
Your Personal Exam Guide

MOHIT CHOUKSEY



$$\frac{1}{2} \times \left(\frac{x}{4}\right) \times \left(\frac{x}{2}\right)$$

$$\left(\frac{x^2}{16}\right) \times 4$$

Q172

$$\frac{x \times x/2}{2}$$

$$\frac{x^2}{4}$$

Q163 $L \uparrow \quad N \downarrow \rightarrow e^x$

80 units
Load \uparrow

100 cycles
 $N \uparrow$

40 units

10,000 $\leftarrow N$


5,000 $\leftarrow N$

$\checkmark 100 \rightarrow 80 \checkmark$
 $\checkmark 10000 \rightarrow 40 \checkmark$
 $\checkmark 5000 \rightarrow \text{---}$

$10,000 \rightarrow 5000$

$$y = k e^{ax}$$

cycle
dependent



independent
variable

$$y = e^{ax}$$

$$y = k a^x$$

$$\frac{k a^{80}}{k a^{40}} = \frac{100}{10000}$$

$$a^{40} = \frac{1}{100}$$




MOHIT CHOUKSEY

(169) $\frac{x}{100}y + \frac{y}{100}x$

$\frac{2xy}{100}$

2% of xy


(157) $(\ln x, y)$



$\tan \theta = -0.02$

$\log_e x = 1$

$x = e^1$



(SIR) $(y - y_1) = m(x - x_1)$



$(x_1, y_1) \rightarrow m$
 \swarrow \nearrow
 $\ln x$ general

$\ln x = .1$


$(y - 0) = m(x - .1)$

$(y - 0) = -.02(x - .1)$

$y = \frac{-2}{100}(x - .1)$

$y = \frac{-2}{100}(\ln 5 - .1)$


(156) $3x + 2y = 14$
 $2x - 3y = 5$



$3x + 2y = 14$
 $x = 0 \quad y = 0 \quad 4.66$
 $y = 7 \quad x = 14/3$
 $x = 4.6$

$2x - 3y = 5$
 $x = 0 \quad y = 0 \quad 2.5$
 $y = -5/3 \quad x = 5/2$

$$\frac{1}{2} \times \frac{14}{3} \times 7 - \frac{1}{2} \left(\frac{14}{3} - \frac{5}{2} \right) \times 4$$



$$\frac{13}{7} \times \frac{6}{8}$$


$$\frac{98}{6} - \frac{13}{12}$$

$$\frac{12 \times 98 - 13 \times 6}{72}$$

$$\frac{1086 - 78}{72}$$

$$= 15.25$$

$$\begin{array}{r} 98 \\ 12 \\ \hline 106 \\ 98 \times \\ \hline 1086 \end{array}$$



ABCD (Trapezium)

$$\Rightarrow \frac{1}{2} (7 + 1) \times 4 = \frac{8}{2} \times 4$$




$$= \frac{32}{2} = 16$$

$$\frac{1}{2} \times 1.5 \times 1 = -.75$$

$$15.25$$


MOHIT CHOUKSEY

150



$A_s + A_R \rightarrow \text{minm.}$

SIR



$\frac{x^2}{x}$

$\frac{2l}{2l^2}$

$4x + 6l = 340$

$[l = \frac{340 - 4x}{6}]$

$A = x^2 + 2l^2$

$A = x^2 + (\frac{340 - 4x}{6})^2$

141

50% ← prone TB → infection

30% ← ~~infected~~ → develops the disease.

70% ✓ (C) ✓

14b

S, M, E, F


$M \rightarrow 2y \rightarrow \frac{1}{2} E$

$S, M \rightarrow 6h$

$E, F \rightarrow 12h$

MOHIT CHOUKSEY

135



$$\frac{PS}{QS} = \frac{3}{1}$$

$$\frac{PU}{UR} = \frac{1}{1}$$

$$\frac{RT}{QT} = \frac{5}{2}$$

$$A_{QTS} = 20 \text{ cm}^2$$

$$\frac{1}{2} \times UT \times RT + \frac{1}{2} \times UT \times TQ + \frac{1}{2} \times PS \times QT$$

$$\frac{1}{2} \times UT (RT + QT) + \frac{1}{2} \times PS (QT)$$

$$\frac{1}{2} \times \frac{UT}{QS} \times QT \left(\frac{RT}{QT} + 1 \right) + \frac{1}{2} \times \frac{PS}{QS} (QT \times QS)$$

$$\frac{1}{2} \times UT \times QT \left(\frac{5}{2} + 1 \right) + \frac{1}{2} \times 3 (QT \times QS)$$

$$\frac{1}{2} \times UT \times QT \left(\frac{7}{2} \right) + 60$$

$$\frac{1}{2} \times \frac{UT}{QS} \times (QT \times QS) \times \frac{7}{2} + 60$$

$$\frac{UT}{QS} \times 35 + 60 = 70 + 60$$

$$\frac{1}{2} \times QT \times QS = 20$$


$$40$$

$$12 = \sqrt{5^2 + 6^2}$$


$$1 - 5^2 =$$

-4

138



132.




$r = 30\text{cm}$

Remaining area = $0.9 \times \pi (30)^2 = \text{lateral surface area of the cone}$

$= \pi R (l)$
 ← slant height (l)

$\Rightarrow R = 27$
 $\Rightarrow r = 30$




$h^2 + 27^2 = 30^2$

$h = \sqrt{30^2 - 27^2}$


$\frac{R}{h} = \frac{27}{13.076}$

$h = 13.076$


*




arc length $\frac{10}{360} (2\pi R)$ area also 10% ↓
 $\frac{10}{360} (\pi R^2)$
 Linear dependency. $\Rightarrow (2\pi R) 0.9 = 2\pi R$



$30 \quad R = 27$



$P(x_1, y_1)$
 $Q(x_2, y_2)$
 $R(x_3, y_3)$




$-4 \leq x \leq 5$
 $-6 \leq y \leq 16$


$\left\{ \begin{array}{l} -4 \leq x \leq 5 \\ 6 \leq y \leq 16 \end{array} \right\}$

$\angle P = 90^\circ$, $PR \parallel x$ axis

$PQR \rightarrow$ integer coordinate



$[{}^{11}C_2 \times {}^{10}C_2] \times 4$



126, 127, 136, ✓

MOHIT CHOUKSEY

120

$$a^2 + b^2 + c^2 = 1$$

$$ab + bc + ca$$

$$(a+b+c)^2 = \underbrace{a^2 + b^2 + c^2}_1 + 2 \underbrace{(ab + bc + ca)}_{1+2}$$

$$1 + 2$$

$$(a+b+c)^2 - (a^2 + b^2 + c^2) = 2(ab + bc + ca)$$

$(-1) + (a+b+c)^2 = 2(ab + bc + ca)_{\min}$
 ↓ for making min this value make = 0
 → +ve/0
 ↳ min $(-\frac{1}{2})$


But

$\left[-\frac{1}{2}, \frac{1}{2} \right]$
 $\left[-\frac{1}{2}, 1 \right]$ in cat

$$(a-b)^2 + (b-c)^2 + (c-a)^2 = 2(a^2 + b^2 + c^2) - 2(ab + bc + ca)$$

$$2(ab + bc + ca)_{\max} = 2(1) - [(a-b)^2 + (b-c)^2 + (c-a)^2]$$

122



- 15 ✓
- 2 x $\frac{6}{3}$ 21 ✓
- 3 x $\frac{3}{1}$ 24 ✓
- 9 15 ✓
- 4 x $\frac{3}{3}$ 41

SIR

- 6 5 ✓ 4
- 7+4 ✓ 7 2+1
- 1+9+2 ✓ 8 1+2+1

118

M → M
P → P
C → C

$$P + m + c = 27/20$$


$$P + m + c = 13/20$$

$$P \times m \times c = 1/10$$

75% → atleast one


50% → atleast two

40% → exactly two



SIR

$$S = 10\% \cdot \frac{1}{10}$$



$$R + B + S = 75$$

\downarrow \downarrow \downarrow
 25 40 10

$m \times p \times c = 1/10$

$$(2) \quad m + p + c = \frac{13}{20} = \frac{65}{100} = 65\% \quad \therefore < 75\%$$

$$m + p + c = \frac{135}{100} = \frac{27}{20}$$


$$R + 2B + 3S$$

$$25 + 2(40) + 3(10) = 135$$

112/113

MOHIT CHOUKSEY

112



113 only read h

rem $\left(\frac{p \times q}{r \times s}\right)$ if $(p \times q) > (r \times s)$

SIR $h = \text{Re}\left(\frac{7 \times 3}{5 \times 2}\right) = \sqrt{1}$



fg(1, 6, 8)

f(1, 4, 6, 8) g(1, 4, 6, 8)






max(p, q, r, s) min(p, q, r, s)

max 8 x 1 = 8

112




MOHIT CHOUKSEY



$$\frac{h}{a} = \sin c$$

$$h = a \sin c$$

$$\frac{1}{2} \times b \times h = \frac{1}{2} \times a b \sin c$$



$$\frac{1}{2} \times q \times PS \sin 60^\circ + \frac{1}{2} \times r \times PS \sin 60^\circ$$


$$\Delta = \frac{1}{2} q r \sin(120^\circ)$$

$$PS(q+r) = qr$$

MOHIT CHOUKSEY

89

Cal	74
Sc	50
Both	106
None	70



$$\left[\frac{74 + 70}{300} \right] \times 100$$

88 D → 10% → T.F.

SIR

$$(100 \times 2) S = 1000$$

$$\begin{array}{r} 100 - 10 \\ - 5 \\ \hline 15 \end{array}$$

1 Ticket → 85/-

$$(85 \times 2) \times 5 = 850$$

76

100B → 4 B

R → 1B → defective
5DB ✓

SIR

T = 100
D = 5
D = 95

$$\frac{f_c}{T_c} = \frac{{}^{95}C_4}{{}^{100}C_4} \quad (.95)^4$$

Total chances

73

population

66Q

[HH] [HT] [TH]

$$\frac{1}{2} \times \frac{1}{2}$$

$$= \frac{1/4}{3/4} = 1/3$$

$$\left[\frac{1}{2} \times 1 + \frac{1}{2} \times \frac{1}{2} \right]$$

MOHIT CHOUKSEY